

LIE ALGEBROID

Lie algebroids were first introduced and studied by J. Pradines [11], following work by C. Ehresmann and P. Libermann on *differentiable groupoids* (later called *Lie groupoids*). Just as Lie algebras are the infinitesimal objects of Lie groups, Lie algebroids are the infinitesimal objects of **Lie groupoids**. They are generalizations of both **Lie algebras** and **tangent vector bundles**. For a comprehensive treatment and lists of references, see [8], [9]. See also [1], [4], [6], [13], [14].

A real *Lie algebroid* $(A, [\cdot, \cdot]_A, q_A)$, is a smooth real vector bundle A over base M , with a real Lie algebra structure $[\cdot, \cdot]_A$ on the vector space $\Gamma(A)$ of the smooth global sections of A , and a morphism of vector bundles $q_A : A \rightarrow TM$, where TM is the tangent bundle of M , called the *anchor*, such that

- $[X, fY]_A = f[X, Y]_A + (q_A(X).f) Y$, for all $X, Y \in \Gamma(A)$ and $f \in C^\infty(M)$,
- q_A defines a Lie algebra homomorphism from the Lie algebra of sections of A , with Lie bracket $[\cdot, \cdot]_A$, into the Lie algebra of vector fields on M .

Complex Lie algebroid structures [1] on complex vector bundles over real bases can be defined similarly, replacing the tangent bundle of the base by the complexified tangent bundle.

The space of sections of a Lie algebroid is a **Lie-Rinehart algebra**, also called a Lie d -ring or a Lie pseudoalgebra. (See [4], [6], [9].) More precisely, it is an (R, \mathcal{A}) -Lie algebra, where R is the field of real (or complex) numbers, and \mathcal{A} is the algebra of functions on the base manifold. In fact, the Lie-Rinehart algebras are the algebraic counterparts of the Lie algebroids, just as the modules over a ring are the algebraic counterparts of the vector bundles.

Examples

1. A Lie algebroid over a one-point set, with the zero anchor, is a Lie algebra.
2. The tangent bundle, TM , of a manifold M , with bracket the Lie bracket of vector fields and with anchor the identity of TM , is a Lie algebroid over M . Any integrable sub-bundle of TM , in particular the

tangent bundle along the leaves of a **foliation**, is also a Lie algebroid.

3. A vector bundle with a smoothly varying Lie algebra structure on the fibers (in particular, a Lie algebra bundle [8]) is a Lie algebroid, with pointwise bracket of sections and zero anchor.

4. If M is a **Poisson manifold** then the cotangent bundle T^*M of M is, in a natural way, a Lie algebroid over M . The anchor is the map $P^\sharp : T^*M \rightarrow TM$ defined by the Poisson bivector P . The Lie bracket $[\cdot, \cdot]_P$ of differential 1-forms satisfies $[df, dg]_P = d\{f, g\}_P$, for any functions $f, g \in C^\infty(M)$, where $\{f, g\}_P = P(df, dg)$ is the Poisson bracket of functions, defined by P . When P is nondegenerate, M is a symplectic manifold and this Lie algebra structure of $\Gamma(T^*M)$ is isomorphic to that of $\Gamma(TM)$. For references to the early occurrences of this bracket, which seems to have first appeared in [3], see [4], [6] and [13]. It was shown in [2] that $[\cdot, \cdot]_P$ is a Lie algebroid bracket on T^*M .

5. The Lie algebroid of a Lie groupoid $(\mathcal{G}, \alpha, \beta)$, where α is the source map and β is the target map [11] [8] [13]. It is defined as the normal bundle along the base of the groupoid, whose sections can be identified with the right-invariant, α -vertical vector fields. The bracket is induced by the Lie bracket of vector fields on the groupoid, and the anchor is $T\beta$.

6. Atiyah sequence. If P is a principal bundle with structure group G , base M and projection p , the G -invariant vector fields on P are the sections of a vector bundle with base M , denoted TP/G , and sometimes called the *Atiyah bundle* of the principal bundle P . This vector bundle is a Lie algebroid, with bracket induced by the Lie bracket of vector fields on P , and with surjective anchor induced by Tp . The kernel of the anchor is the adjoint bundle, $(P \times \mathfrak{g})/G$. Splittings of the anchor are **connections** on P . The Atiyah bundle of P is the Lie algebroid of the Ehresmann gauge groupoid $(P \times P)/G$. If P is the frame bundle of a vector bundle E , then the sections of the Atiyah bundle of P are the covariant differential operators on E , in the sense of [8].

7. Other examples are the trivial Lie algebroids $TM \times \mathfrak{g}$, the transformation Lie algebroids $M \times \mathfrak{g} \rightarrow M$, where Lie algebra \mathfrak{g} acts on manifold M , the deformation Lie algebroid $A \times \mathbb{R}$ of a Lie algebroid A ,

where $A \times \{\hbar\}$, for $\hbar \neq 0$, is isomorphic to A , and $A \times \{0\}$ is isomorphic to vector bundle A with the abelian Lie algebroid structure (zero bracket and zero anchor), the prolongation Lie algebroids of a Lie algebroid, etc.

de Rham differential. Given any Lie algebroid A , a differential d_A is defined on the graded algebra of sections of the exterior algebra of the dual vector bundle, $\Gamma(\wedge A^*)$, called the *de Rham differential* of A . Then $\Gamma(\wedge A^*)$ can be considered as the algebra of functions on a **supermanifold**, d_A being an odd vector field with square zero [12].

If A is a Lie algebra \mathfrak{g} , then d_A is the Chevalley-Eilenberg cohomology operator on $\wedge(\mathfrak{g}^*)$.

If $A = TM$, then d_A is the usual de Rham differential on forms.

If $A = T^*M$ is the cotangent bundle of a Poisson manifold, then d_A is the Lichnerowicz-Poisson differential $[P, \cdot]_A$ on fields of multivectors on M .

Schouten algebra. Given any Lie algebroid A , on the graded algebra of sections of the exterior algebra of vector bundle A , $\Gamma(\wedge A)$, there is a *Gerstenhaber algebra* structure (see **Poisson algebra**), denoted by $[\cdot, \cdot]_A$. With this graded Lie bracket, $\Gamma(\wedge A)$ is called the *Schouten algebra* of A .

If A is a Lie algebra \mathfrak{g} , then $[\cdot, \cdot]_A$ is the *algebraic Schouten bracket* on $\wedge \mathfrak{g}$.

If $A = TM$, then $[\cdot, \cdot]_A$ is the usual Schouten bracket of fields of multivectors on M .

If $A = T^*M$ is the cotangent bundle of a Poisson manifold, then $[\cdot, \cdot]_A$ is the Koszul bracket [7] [13] [5] of differential forms.

Morphisms of Lie algebroids and the linear Poisson structure on the dual. A base-preserving morphism from Lie algebroid A_1 to Lie algebroid A_2 , over the same base M , is a base-preserving vector-bundle morphism, $\mu : A_1 \rightarrow A_2$, such that $q_{A_2} \circ \mu = q_{A_1}$, inducing a Lie-algebra morphism from $\Gamma(A_1)$ to $\Gamma(A_2)$.

If A is a Lie algebroid, the dual vector bundle A^* is a *Poisson vector bundle*. This means that the total space of A^* has a Poisson structure such that the Poisson bracket of two functions which are linear on the fibers is linear on the fibers. A base-preserving morphism from vector bundle A_1 to vector bundle A_2 is a morphism of Lie algebroids if and only if its transpose is a Poisson morphism.

Lie bialgebroids [10] [5] are pairs of Lie algebroids (A, A^*) in duality satisfying the compatibility condition that d_{A^*} be a derivation of the graded Lie bracket $[\cdot, \cdot]_A$. They generalize the Lie bialgebras in the sense of V. G. Drinfel'd (see **quantum groups** and **Poisson Lie groups**) and also the pair (TM, T^*M) , where M is a Poisson manifold.

There is no analogue to Lie's third theorem in the case of Lie algebroids, since not every Lie algebroid can be integrated to a global Lie groupoid, although there are local versions of this result. (See [8], [1].)

References

- [1] CANNAS DA SILVA, A., and WEINSTEIN, A.: *Geometric Models for Noncommutative Algebras*, Berkeley Math. Lect. Notes **10**, Amer. Math. Soc., 1999.
- [2] COSTE, A., DAZORD, P., and WEINSTEIN, A.: 'Groupoïdes symplectiques', *Publications du Département de Mathématiques, Université Claude Bernard-Lyon I*, **2A** (1987), 1-62.
- [3] FUCHSSTEINER, B.: 'The Lie algebra structure of degenerate Hamiltonian and bi-Hamiltonian systems', *Prog. Theor. Phys.* **68** (1982), 1082-1104.
- [4] HUEBSCHMANN, J.: 'Poisson cohomology and quantization', *J. Reine Angew. Math.* **408** (1990), 57-113.
- [5] KOSMANN-SCHWARZBACH, Y.: 'Exact Gerstenhaber algebras and Lie bialgebroids', *Acta Appl. Math.* **41** (1995), 153-165.
- [6] KOSMANN-SCHWARZBACH, Y., and MAGRI, F.: 'Poisson-Nijenhuis structures', *Ann. Inst. Henri Poincaré, Phys. Théor.* **53** (1990), 35-81.
- [7] KOSZUL, J.-L.: 'Crochet de Schouten-Nijenhuis et cohomologie', *Astérisque*, hors série (1985), 257-271.
- [8] MACKENZIE, K.: *Lie Groupoids and Lie Algebroids in Differential Geometry*, Cambridge Univ. Press, 1987.
- [9] MACKENZIE, K.: 'Lie algebroids and Lie pseudoalgebras', *Bull. London Math. Soc.* **27** (1995), 97-147.
- [10] MACKENZIE, K., and XU, P.: 'Lie bialgebroids and Poisson groupoids', *Duke Math. J.* **73** (1994), 415-452.
- [11] PRADINES, J.: 'Théorie de Lie pour les groupoïdes différentiables. Calcul différentiel dans la catégorie des groupoïdes infinitésimaux', *Comptes rendus Acad. Sci. Paris* **264 A** (1967), 245-248.

- [12] VAINTROB, A.: ‘Lie algebroids and homological vector fields’, *Russ. Math. Surv.* **52** (1997), 428-429.
- [13] VAISMAN, I.: *Lectures on the Geometry of Poisson Manifolds*, Birkhäuser, 1994.
- [14] WEINSTEIN, A.: Poisson Geometry, *Diff. Geom. Appl.* **9** (1998), 213-238.

Yvette Kosmann-Schwarzbach

AMS Classification: 58H05, 58A99, 17B66, 17B70, 53C15, 14F40, 22A22.

to appear in *Encyclopaedia of Mathematics*
(Supplement II), Kluwer