

Commuting Hamiltonians and Hamilton-Jacobi multi-time equations

FRANCO CARDIN ¹ CLAUDE VITERBO ²

¹ Dipartimento di Matematica Pura ed Applicata
via Trieste 63 - 35121 Padova, Italia
`cardin@math.unipd.it`

² Centre de Mathématiques Laurent Schwartz
UMR 7640 du CNRS
École Polytechnique - 91128 Palaiseau, France
`viterbo@math.polytechnique.fr`

September 7, 2007

Abstract

The aim of this paper is twofold: first of all, we show that the C^0 limit of a pair of commuting Hamiltonians commute. This means on one hand that if the limit of the Hamiltonians is smooth, the Poisson bracket of their limit still vanishes, and on the other hand that we may define “commutation” for C^0 functions.

The second part of the paper deals with solving “multi-time” Hamilton-Jacobi equations using variational solutions. This extends the work of Barles and Tourin in the viscosity case to include the case of C^0 Hamiltonians, and removes their convexity assumption, provided we are in the framework of “variational solutions”.

Hamilton's variation principle can be shown to correspond to Fermat's *Principle* for a wave propagation in configuration space (q -space), and the Hamilton-Jacobi equation expresses Huygens' *Principle* for this wave propagation. Unfortunately this powerful and momentous conception of Hamilton, is deprived, in most modern reproductions, of its beautiful raiment as a superfluous accessory, in favour of a more colourless representation of the analytical correspondence.

E. Schrödinger, Quantization as a Problem of eigenvalues
(Part II), *Annalen der Physik*, 1926

1 Introduction

The problem of finding solutions of multi-time Hamilton-Jacobi equations, by which one usually means equations of the following type, where x is in \mathbb{R}^n and t_j in \mathbb{R}

$$(MHJ) \quad \begin{cases} \frac{\partial}{\partial t_1} u(t_1, \dots, t_d, x) + H_1(t_1, \dots, t_d, x, \frac{\partial}{\partial x} u(t_1, \dots, t_d, x)) = 0 \\ \vdots \\ \frac{\partial}{\partial t_d} u(t_1, \dots, t_d, x) + H_d(t_1, \dots, t_d, x, \frac{\partial}{\partial x} u(t_1, \dots, t_d, x)) = 0 \end{cases}$$

with initial condition

$$u(0, \dots, 0, x) = f(x)$$

has been initiated by Rochet in relation with some questions in economy, then by Lions-Rochet and studied more recently by Barles-Tourin and Motta-Rampazzo ([Rochet, Lions-Rochet, Barles-Tourin, Motta-Rampazzo]). Such a system of equations is well-known to be overdetermined, and in order to have a solution, we need the Hamiltonians to commute in a suitable sense. This is already obvious when applying the method of characteristics. Besides a suitable commutation condition, we need to address the question of the type of solution one is looking for. For first order equations, it is well known, and was proved for more general equations by Dacorogna and Marcellini [Dacorogna-Marcellini], that there are plenty of C^0 solutions for such equations¹. We then need to select a particular “class” of solution, deemed to be the best suited to our problem. A classical choice is to look for viscosity solutions, which are the “right” solutions for optimal control and this is the type of solution considered in the above papers (except for [Dacorogna-Marcellini]).

In Barles and Tourin’s paper, the existence of a viscosity solution is proved under the assumption that one of the Hamiltonians is coercive with controlled growth²

(see [Barles-Tourin], page 1526, conditions (H1),(H2)), and more importantly that H_1, H_2 satisfy the following conditions

- (a). independent of (t_1, \dots, t_d)
- (b). convex in p
- (c). of class C^1 and satisfying the commuting condition $\{H_j, H_k\} = 0$

In fact the third condition can be weakened, as Barles and Tourin point out, to assume that the H_j are C^0 and there are sequences of Hamiltonians H_j^ν of class C^1 such that

¹By C^0 solution we mean C^1 almost everywhere, and satisfying the equation a.e.

²Condition (H1) in [Barles-Tourin] states that for each R there exists K_R such that $|H_1(x, p)| \leq K_R$ and $|\frac{\partial H_1}{\partial p}(x, p)| \leq K_R(1 + |x|)$ in $\mathbb{R}^n \times B(0, R)$. Moreover the authors assume $du_0(x)$ to be bounded. These conditions are stronger than the assumptions we need here. See appendix B for more details.

(a). $\lim_{\nu \rightarrow \infty} H_j^\nu = H_j$ in the C^0 topology

(b). $\{H_j^\nu, H_l^\nu\} = 0$

In other words, the H_j are limits of commuting Hamiltonians.

However such an assumption is quite unpractical since it is already difficult to write two commuting C^1 Hamiltonians as nontrivial limits of commuting Hamiltonians.

The present paper has several goals.

First we solve the multi-time Hamilton-Jacobi equation in the framework of “variational solutions” defined by Sikorav, Chaperon and the second author in [Sikorav 1, Chaperon, Viterbo-Ottolenghi] (see definition 2.1). According to a result by Zhukovskaia (cf [Zhukovskaya]), if the Hamiltonian is convex in p , the variational solution must coincide with the viscosity solution defined by Crandall and Lions (in [Crandall-Lions], see also our definition 2.11), so that our results extend those of Barles and Tourin. However in general these two solutions do not coincide (see an example in [Viterbo 2]). For variational solutions, we prove that we only need the H_j to be C^0 , and condition (b) is then replaced by a “commutation” condition best expressed in terms of symplectic invariants, refining the following

Definition 1.1. Let H, K be two autonomous C^0 Hamiltonians. We shall say that H and K **C^0 -commute**, if and only if there are sequences H_ν, K_ν of C^1 Hamiltonians such that, all limits being for the C^0 topology, we have:

(a). $\lim_{\nu \rightarrow \infty} H_\nu = H, \lim_{\nu \rightarrow \infty} K_\nu = K.$

(b). $\lim_{\nu \rightarrow \infty} \{H_\nu, K_\nu\} = 0.$

A similar definition is given in section 3 for H, K time-dependent and in appendix C for the case of equations depending on the function.

Postponing to the next Section the detailed geometrical setting for our Hamiltonians, the two main theorems of this paper are:

Theorem 1.2. *If two $C^{1,1}$ Hamiltonians C^0 -commute, then they commute in the usual sense (i.e. their Poisson bracket vanishes).*

Here $C^{1,1}$ means differentiable with Lipschitz differential in the variables (x, p) . The above theorem tells us that our definition of C^0 -commutation coincides with the classical one for smooth Hamiltonians. Note that this may be extended to the time-dependent setting as we shall see in section 3.1.

The above theorem sounds like a generalization of Eliashberg-Gromov’s theorem ([Gromov], [Eliashberg], [Ekeland-Hofer]) on the C^0 closure of the group of symplectic diffeomorphism, according to which the set of $2n$ -tuples of functions on \mathbb{R}^{2n} , $(f_1, \dots, f_n, g_1, \dots, g_n)$ such that

$$\{f_i, f_j\} = \{g_i, g_j\} = 0, \{f_i, g_j\} = \delta_i^j$$

is closed³. We refer to [Humilière 2] for an approach along these lines of the Gromov-Eliashberg theorem. We also refer to improvements of the above result from a quantitative point of view due to [Entov-Polterovich-Zapolsky] using quasi-states.

This has been extended by V. Humilière to other relations derived from so-called quasi-representations of finite dimensional Lie groups in the Poisson algebra. For example the Heisenberg relation $\{f, g\} = h; \{f, h\} = \{g, h\} = 0$ is also C^0 closed (see [Humilière 2]).

Theorem 1.3. *Assume the Hamiltonians $H_1(t_1, \dots, t_d, x, p), \dots, H_d(t_1, \dots, t_d, x, p)$ on $T\mathbb{R}^n$ satisfy the following conditions*

- (a). *they are locally Lipschitz in (x, p) and their Lipschitz constant on the ball of radius r has at most linear growth in r .*
- (b). *their support has an x -projection contained in a compact set.*
- (c). *They C^0 -commute.*

Then equation (MHJ) has a unique solution which is a variational solution of each individual equation. If all the Hamiltonians H_j 's are convex in p , then u is a viscosity solution of each individual equation.

Remarks 1.4. (a). We refer to definition 2.1 and 2.11 for the meaning of variational and viscosity solution.

- (b). The growth condition is only needed to guarantee the existence of the flow for the approximating Hamiltonians. Since if dH has linear growth, we may approximate it by smooth Hamiltonians H^ν such that the vector field X_{H^ν} is complete⁴, this condition is sufficient. In fact, it is enough to assume there are constants A, B such that $|X_H(q, p)| \leq A(|q| + |p|) + B$, i.e. the Lipschitz norm of H on a ball grows at most linearly with the radius. Note also that in some cases, we may guarantee existence of the flow of the X_{H^ν} for other reasons. For example when H is autonomous, and proper, since the H^ν will satisfy the same assumption, and conservation of energy implies that the flow remains in a compact set, the flow of X_H is thus defined for all times.

Remark 1.5. The paper is not supposed to be written for specialists in symplectic topology, although a certain familiarity with the basic constructions of [Viterbo 1] and [Viterbo-Ottolenghi] is recommended. Appendices A and B are of a more symplectic flavor and really address the question of Hamilton-Jacobi equations from a

³To be rigorous, Gromov and Eliashberg need the map $(x, y) \longrightarrow (f_1(x, y), \dots, f_n(x, y), g_1(x, y), \dots, g_n(x, y))$ to be bijective.

⁴that is, the flow φ^t is defined for all t in \mathbb{R}

symplectic topology viewpoint. In particular Appendix B addresses the question of the growth conditions one must impose on the Hamiltonian and the initial condition from a purely geometric point of view, while Appendix C extends the main theorem for equations depending on the unknown function.

1.1 Organization of the paper

Section 2 is devoted to a summary of the applications of Generating function theory to symplectic topology, in a slightly modified version with respect to [Viterbo 1]. We also state the main properties of variational solutions as in [Viterbo-Ottolenghi].

Section 3 deals with the proof of theorem 1.2. The proof is based on continuity properties of the symplectic norm c defined in Section 2. We prove that if $\{H_n, K_n\}$ are C^0 , small, the flows φ_n^t, ψ_n^s of H_n and K_n have the following properties:

on one hand $t \longrightarrow \varphi_n^t \psi_n^s \varphi_n^{-t} \psi_n^{-s}$ is generated by a C^0 small Hamiltonian, and the properties of c established in the previous section imply that $c(\varphi_n^t \psi_n^s \varphi_n^{-t} \psi_n^{-s})$ is small. On the other hand, if H_n goes to H with flow φ^t and K_n goes to K with flow ψ^t , $(\varphi_n^t \psi_n^s \varphi_n^{-t} \psi_n^{-s})$ goes to $\varphi^t \psi^s \varphi^{-t} \psi^{-s}$. Uniqueness of limits and the fact that c only vanishes on the identity implies that $\varphi^t \psi^s \varphi^{-t} \psi^{-s} = \text{Id}$ for all s, t , hence H and K commute.

In section 4 we first show how multi-time equations have natural variational solutions, provided the Hamiltonians commute. We then address a number of technical questions, replacing the invariants of section 2 by their stabilization.

Section 5 eventually completes the proof of theorem 1.3. It is sufficient to deal with equation (MHJ) in the case of two Hamiltonians. Assume the two Hamiltonians are such that $\{H_1, H_2\}$ is C^0 small. We then construct two Lagrangians, $L_{1,2}$ and $L_{2,1}$ obtained by “solving” the first of the two equations (for $t_2 = 0$) and then the second, and vice versa. We must then prove that the two Lagrangians $L_{1,2}$ and $L_{2,1}$ are close with respect to the γ distance defined in [Viterbo 1] and also in subsection 2.1 (this is not so with respect to the C^0 distance). Once this is granted, it implies that the associated function $u_{1,2}$ and $u_{2,1}$ are C^0 close. The proof of the theorem is now obtained by limiting arguments.

Appendix A gives the proof of some technical results. Appendix B extends the scope of the main theorems to the case of a non compact support. Appendix C,D,E give some complements on equations involving the unknown function, the geometric theory of Hamilton-Jacobi equations associated to coisotropic manifolds and historical comments.

1.2 Acknowledgements

The authors warmly thank Franco Rampazzo for attracting their attention to this problem during a conference in Cortona, for communicating his lecture notes on his

work [Motta-Rampazzo] and for many interesting discussions. We also would like to thank F.Camilli, I.Capuzzo Dolcetta and A.Siconolfi for the superb organization of the Cortona conference. Even though it is probably not related, we also mention the paper [Rampazzo-Sussmann] on commutation of Lipschitz vector fields.

2 Preliminary material

2.1 Generating functions and variational solutions of Hamilton-Jacobi equations

We shall here assume that N is a connected manifold without boundary, and either compact or that all Hamiltonians are compact supported, as in [Viterbo 1]. However, we shall explain in Appendix B, how our results extend to non-compact situations, provided we have some estimate on the growth of the Hamiltonians.

Let T^*N be the cotangent bundle of the manifold N endowed with the canonical symplectic structure $\sigma = \sum_{j=1}^n dp_j \wedge dx^j$. To any Hamiltonian $H(t, z)$ (where $z = (x, p)$) on $\mathbb{R} \times T^*N$ we associate the time-dependent vector field X_H defined by

$$\sigma(X_H, \xi) = -d_z H(t, z)\xi$$

and the corresponding Hamiltonian flow $\varphi_{t_0}^t$ defined by

$$\begin{cases} \frac{d}{dt}\varphi_{t_0}^t = X_H(t, \varphi_{t_0}^t) \\ \varphi_{t_0}^{t_0} = Id \end{cases}$$

Let L be a Lagrangian submanifold of T^*N obtained from the zero section $0_N = \{(x, 0) \in T^*N \mid x \in N\}$ by the Hamiltonian isotopy $\varphi_{t_0}^t$. We shall always assume that $\varphi_{t_0}^t$ is well defined on 0_N for all t . Then according to [Sikorav-2] (relying on joint work with Laudenbach in [Laudenbach-Sikorav]), there exists a generating function quadratic at infinity for L i.e. there exists a smooth function $S : N \times \mathbb{R}^k \rightarrow \mathbb{R}$ such that

- (a). $(x, \xi) \mapsto \frac{\partial S}{\partial \xi}(x, \xi)$ has 0 as a regular value (a Morse family in the terminology introduced by A. Weinstein in [Weinstein])
- (b). $S(x, \xi) = Q(\xi)$ for ξ large enough, where Q is a non-degenerate quadratic form⁵.

⁵This is sometimes conveniently replaced by the condition $|S(x, \cdot) - Q(\cdot)|_{C^1} < +\infty$. We shall use these conditions interchangeably in the rest of the paper. It is easy to prove that existence of a generating function of one kind is equivalent to the existence of a generating function of the other kind, see [Brunella].

(c).

$$L = \{(x, \frac{\partial S}{\partial x}(x, \xi)) \mid \frac{\partial S}{\partial \xi}(x, \xi) = 0\}$$

In particular the critical points of S are in one to one correspondence with the points of $L \cap 0_N$.

In the rest of the paper, we shall shorten the expression “generating function quadratic at infinity” by “GFQI”.

Let $S^\lambda = \{(x, \xi) \mid S(x, \xi) \leq \lambda\}$, E^\pm be the positive and negative eigenspaces of Q , and D^\pm be large discs in E^\pm . Since for c large enough $S^{\pm c} = N \times Q^{\pm c}$ we have

$$H^*(S^c, S^{-c}) = H^*(N \times Q^c, N \times Q^{-c}) = H^*(N) \otimes H^*(D^-, \partial D^-)$$

so that, to each cohomology class $\alpha \in H^*(N)$ we may associate a class, image of α by the Künneth isomorphism, denoted by $T\alpha$. To the class $T\alpha$ in $H^*(S^c, S^{-c})$, we may associate a minimax critical level

$$c(\alpha, S) = \inf\{\lambda \mid T\alpha \notin \text{Ker}(H^*(S^c, S^{-c}) \rightarrow H^*(S^\lambda, S^{-c}))\}$$

Now it is proved in [Theret 2] and [Viterbo 1] that given L , S is essentially unique up to adding a constant, and more precisely, up to a global shift, the numbers $c(\alpha, S)$ depend only on L , not on S , and they are thus denoted by $c(\alpha, L)$. Moreover, denoting by

$$\gamma(L) = c(\mu, L) - c(1, L)$$

where $1 \in H^0(N)$, $\mu \in H^n(N)$ are generators. We know that $\gamma(L)$ is well defined and vanishes if and only if $L = 0_N$ (see for instance [Viterbo 1]).

Moreover let $S_x(\xi) = S(x, \xi)$ be the restriction of S to the fiber over x . We can look for a minimax as above for the function S_x . Since the cohomology of the point is one dimensional, denoting its generator by 1_x , we set

Definition 2.1. The continuous function

$$u_L(x) = c(1_x, S_x)$$

is called **the variational solution** of the equation

$$(x, du(x)) \in L.$$

In particular if $L \subset H^{-1}(0)$ u_L is a **variational solution** of the Hamilton-Jacobi equation

$$H(x, du(x)) = 0$$

It has been proved by Sikorav and Chaperon (see [Chaperon, Sikorav 1] and also [Viterbo-Ottolenghi]), that such a function is indeed a solution of the *Lagrangian Hamilton-Jacobi equation*, that is

$$(x, du_L(x)) \text{ is in } L \text{ for almost all } x \text{ in } N$$

When $L \subset H^{-1}(0)$, we have a solution of the classical Hamilton-Jacobi equation

$$H(x, du_L(x)) = 0 \text{ for almost all } x \text{ in } N$$

Of course, for any constant c , $u_L(x) + c$ is also a solution. For evolution equations, that is

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) = 0 \\ u(0, x) = f(x) \end{cases}$$

the construction of variational solutions for a single equation can be rephrased as follows. Let Λ_0 be a Lagrangian submanifold of T^*N , and $H(t, z)$ be a time-dependent Hamiltonian. We consider $\tilde{\Lambda}_{0,H} = \{(0, -H(0, z), z) \mid z \in \Lambda_0\}$, and

$$L = \bigcup_{t \in \mathbb{R}} \Phi^t(\Lambda_{0,H}) \subset T^*(\mathbb{R} \times N)$$

In [Viterbo-Ottolenghi], it is also proved that for evolution equations the definition of variational solutions extends to C^0 Hamiltonians: indeed if H^ν tends to H , then the solution u^ν converges to u . This follows from the property

$$\|u_H - u_K\|_{C^0([0,T] \times \mathbb{R}^n)} \leq T \|H - K\|_{C^0([0,T] \times \mathbb{R}^{2n})}$$

which in turn follows from Proposition 2.6 (see [Viterbo-Ottolenghi, Viterbo 2]). Note that in the framework of viscosity solutions, this property is called **stability**. A priori, even though u is Lipschitz -hence according to Rademacher theorem is almost everywhere differentiable- we do not claim that u satisfies the Hamilton-Jacobi equation almost everywhere⁶.

2.2 Capacities for Hamiltonian flows

Let L_1, L_2 be Lagrangian submanifolds generated by S_1, S_2 . We may define $c(\alpha, S_1 - S_2)$, and as this does not depend on the choice of S_1, S_2 but only on L_1, L_2 we denote it by abuse of language $c(\alpha, L_1 - L_2)$, even though it is not really determined by the set

$$L_1 - L_2 = \{(x, p_1 - p_2) \mid (x, p_1) \in L_1, (x, p_2) \in L_2\}$$

⁶This is the case for viscosity solutions, but is unknown for variational solutions.

but depends on both L_1 and L_2 .

We denote by $\gamma(L_1 - L_2)$ the difference $c(\mu, L_1 - L_2) - c(1, L_1 - L_2)$. This is non-negative according to [Viterbo 1] and vanishes if and only if $L_1 = L_2$.

Now to a compact supported symplectic isotopy, denoted by ψ , we may associate a symplectic invariant as follows:

Definition 2.2. Let \mathcal{L} be the set of Lagrangian submanifolds Hamiltonianly isotopic to the zero section, $\mathcal{H}(T^*N)$ be the set of smooth time-dependent Hamiltonians on $\mathbb{R} \times T^*N$, $\mathcal{DH}(T^*N)$ be the set of time one maps of such Hamiltonians. We shall use the notation $\mathcal{H}, \mathcal{DH}$ if there is no ambiguity. We set for $\psi \in \mathcal{DH}$,

$$\tilde{\gamma}(\psi) = \sup\{\gamma(\psi(L) - L) \mid L \in \mathcal{L}\}$$

We shall need the following

Proposition 2.3. (a).

$$\tilde{\gamma}(\psi) \geq 0$$

and $\tilde{\gamma}(\psi) = 0$ if and only if $\psi = \text{Id}$

(b).

$$\tilde{\gamma}(\psi^{-1}) = \tilde{\gamma}(\psi)$$

(c). (triangle inequality)

$$\tilde{\gamma}(\psi\varphi) \leq \tilde{\gamma}(\psi) + \tilde{\gamma}(\varphi)$$

(d). (invariance by conjugation)

$$\tilde{\gamma}(\varphi\psi\varphi^{-1}) = \tilde{\gamma}(\psi)$$

Proof. The proof of this proposition is postponed to Appendix A □

Definition 2.4. We shall say that the sequence φ_n in \mathcal{DH} *c-converges* to φ if and only if

$$\lim_{n \rightarrow \infty} \tilde{\gamma}(\varphi_n^{-1}\varphi) = 0$$

We shall use the notation

$$\varphi_n \xrightarrow{c} \varphi$$

for *c-convergence*⁷.

⁷*c* stands for capacity, see [Ekeland-Hofer].

Remark 2.5. Since our invariant is called γ , we should talk about γ -convergence. In fact our c -convergence is indeed related to Γ -convergence in the calculus of variations, but we want to avoid any confusion here.

We also need the following estimate:

Proposition 2.6. *Assume $\bar{\psi}$ is the time one map of the Hamiltonian $H(t, z)$. Then we have*

$$\tilde{\gamma}(\psi) \leq \|H\|_{C^0} := \sup_{(t,z) \in [0,1] \times \mathbb{R}^{2n}} H(t, z) - \inf_{(t,z) \in [0,1] \times \mathbb{R}^{2n}} H(t, z)$$

As a consequence, if φ_n and φ are generated by H_n and H , and if $H_n \rightarrow H$ in the C^0 topology, then φ_n c -converges to φ .

Proof. We first prove that if H_0, H_1 are Hamiltonians and φ_0, φ_1 are their flows, we may normalize S_0, S_1 generating functions of $\varphi_0(L), \varphi_1(L)$ so that their critical values are those of

$$\mathcal{S}_j(x(1); \gamma, \xi) = S(x(0), \xi) + \int_0^1 p(t)\dot{x}(t) + H_j(t, x(t), p(t))dt$$

with respect to the (infinite) auxiliary parameters γ, ξ , and where S is generating function quadratic at infinity for L , and $\gamma = (x, p) : [0, 1] \rightarrow T^*N$. Indeed we have

$$\begin{aligned} DS_j(x(1); \gamma, \xi) (\delta x(1), \delta x(0), \delta \gamma, \delta \xi) = \\ p(1)\delta x(1) + \frac{\partial S}{\partial \xi}(x(0), \xi)\delta \xi - \left(p(0) - \frac{\partial S}{\partial x}(x(0), \xi) \right) \delta x(0) + \\ \int_0^1 \left[\left(\dot{x}(t) + \frac{\partial H_j}{\partial p}(x(t), p(t)) \right) \delta p(t) - \left(\dot{p}(t) - \frac{\partial H_j}{\partial x}(x(t), p(t)) \right) \delta x(t) \right] dt \end{aligned}$$

According to [Viterbo 1], [Theret 2] generating functions associated to a Lagrangian are “essentially unique”, up to a constant. Thus the critical values of two such functions differ by a global translation. Since $\mathcal{S}_j(x(1); \gamma, \xi)$ is formally a generating function, and in particular has critical values coinciding up to translation with those of any other generating function, we may use its critical values to normalize the S_j (i.e. we replace S_j by $S_j + c_j$ so that the critical values of $S_j + c_j$ and \mathcal{S}_j coincide).

In particular we claim that if $H_0 \leq H_1$ we have $c(\alpha, S_0) \leq c(\alpha, S_1)$.

For this we argue as in the proof of proposition 4.6 from [Viterbo 1]. We consider the linear interpolation $H_\lambda(t, x, p) = (1 - \lambda)H_0(t, x, p) + \lambda H_1(t, x, p)$. Let φ_λ^t be the flow of H_λ and φ_λ be the time one map. The associated generating function $S_\lambda(x, \xi)$

of $\varphi_\lambda(L)$ is normalized as above. Now for a critical point of S_λ , we get an intersection point $(x_\lambda, 0) = (x_\lambda(1), p_\lambda(1))$ of $\varphi_\lambda(L) \cap 0_N$ and the critical values are those of

$$\mathcal{S}_\lambda(x(1); \gamma, \xi) = S(x(0), \xi) + \int_0^1 [p(t)\dot{x}(t) + H_\lambda(t, x(t), p(t))]dt$$

corresponding to critical points of the form $(x_\lambda(1), \gamma_\lambda, \xi_\lambda)$ where

$$\gamma_\lambda(t) = (x_\lambda(t), p_\lambda(t)) = \varphi_\lambda^t(x_\lambda(0), p_\lambda(0))$$

and

$$\frac{\partial}{\partial \xi} S(x_\lambda(0), \xi_\lambda) = 0$$

Using the fact that generically $x_\lambda = x_\lambda(1)$ is piecewise C^1 , and $S(x_\lambda, \xi_\lambda)$ is continuous, it is enough to know that for all but a finite set of values of λ we may write

$$\frac{d}{d\lambda} S_\lambda(x_\lambda, \xi_\lambda) = \int_0^1 \frac{d}{d\lambda} H_\lambda(t, x_\lambda(t), p_\lambda(t))$$

and this quantity is positive if $H_0 \leq H_1$.

Now $\|H\|_{C^0} = C$ means that $a \leq H \leq b$ with $b - a \leq C$. Then, since for the constant Hamiltonian $h_a(x) = a$,

$$c(\mu, S_a) = c(1, S_a) = a$$

(again because of the above normalization) and we get $a \leq c(1, S_H) \leq c(\mu, S_H) \leq b$ and we have $\gamma(S_H) \leq b - a = \|H\|_{C^0}$ \square

Remark 2.7. As the referee pointed out, A. Weinstein noticed long ago that the action functional is a "generating function" in some generalized sense. Taking some finite dimensional reduction of this, one can associate to H a GFQI S_H such that

$$\|S_H - S_K\| \leq \|H - K\|.$$

Corollary 2.8. *Assume $H_\nu \rightarrow H$ in the C^0 topology, where H is in $C^{1,1}$, (i.e. it has Lipschitz derivatives), and φ_ν^t, φ^t are the flows of H_ν and H . Then for all t , φ_ν^t c -converges to φ^t : $\varphi_\nu^t \xrightarrow{c} \varphi^t$.*

Proof. Indeed $\varphi^{-t}\varphi_\nu^t$ is the flow of $H_\nu(t, \varphi^t(z)) - H(t, \varphi^t(z))$, and clearly if $H_\nu \rightarrow H$ in the C^0 topology, this quantity goes to zero, hence $\tilde{\gamma}(\varphi^{-t}\varphi_\nu^t)$ goes to zero, which is equivalent to $\varphi_\nu^t \xrightarrow{c} \varphi^t$. \square

Remark 2.9. Given φ the time one map of a symplectomorphism of the symplectic manifold M , we may define its **Hofer norm** as follows ([Hofer]) Let $\mathcal{H}(\varphi)$ be the set of (time-dependent) Hamiltonians on M such that the time one flow associated to H is φ .

Then

$$\|\varphi\| = \inf \left\{ \int_0^1 [\max_x H(t, x) - \min_x H(t, x)] dt \mid H \in \mathcal{H}(\varphi) \right\}$$

In fact, the proofs of both proposition 2.6 and corollary 2.7 show more than stated: we prove that the identity map from $(\mathcal{H}, \|\bullet\|)$ to $(\mathcal{H}, \tilde{\gamma})$ is a contraction, or in less pedantic terms, that

$$\tilde{\gamma}(\varphi) \leq \|\varphi\|$$

This allows to set the following definition, as in [Humilière]:

Definition 2.10. ([Humilière]) We define $\mathfrak{H}_\gamma(M)$ the completion of $\mathcal{H}(M)$ for the metric γ , and $\mathfrak{H}\mathcal{D}_\gamma(M)$ the completion of $\mathcal{D}\mathcal{H}(M)$ for the metric $\tilde{\gamma}$.

Let us point out that the “time one flow” map $\mathcal{H}(M) \longrightarrow \mathcal{D}\mathcal{H}(M)$ extends to a continuous map $\mathfrak{H}_\gamma(M) \longrightarrow \mathfrak{H}\mathcal{D}_\gamma(M)$. According to Proposition 2.6, we have a continuous map of $C^0(\mathbb{R} \times M)$ into $\mathfrak{H}_\gamma(M)$. Moreover, according to a theorem of V. Humilière ([Humilière]), Hamiltonians with some controlled singularities also live in $\mathfrak{H}_\gamma(M)$.

2.3 Viscosity solutions

The only fact the reader needs to know here about viscosity solutions is Zhukovskaia’s theorem. For his convenience, we repeat the definition of viscosity solutions in the framework of evolution equations

Definition 2.11. A viscosity subsolution (resp. supersolution) of the Hamilton-Jacobi equation

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) = 0 \\ u(0, x) = f(x) \end{cases}$$

is a function u satisfying the initial condition and such that if $\varphi(t, x)$ is a function such that $u(t, x) - \varphi(t, x)$ has a local maximum (resp. minimum) at (t_0, x_0) we have

$$\frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) \leq 0 \quad (\text{resp. } \geq 0)$$

Moreover u is a viscosity solution if and only if it is both a viscosity subsolution and a viscosity supersolution.

We now have

Theorem 2.12 (Zhukovskaia's theorem). (*[Zhukovskaya], [Bernardi-Cardin]*) *If H is convex in p then a function $u(t, x)$ is a viscosity solution of the Hamilton-Jacobi equation*

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) = 0 \\ u(0, x) = f(x) \end{cases}$$

if and only if it is a variational solution.

Note that an example was given in [Viterbo-Ottolenghi] showing that this theorem fails if we remove the convexity assumption.

3 Commuting autonomous Hamiltonians and the proof of theorem 1.2

Let $f : P_1 \rightarrow P_2$ be a diffeomorphism between two manifolds, and X be a vector field on M_1 , then the *push-forward* of X , f_*X is defined as the vector field: $f_*X(y) = df(f^{-1}(y))X(f^{-1}(y))$. Its main property is that if φ^t is the flow of X , then

$$\frac{d}{dt}(f \circ \varphi^t)(x) = (f_*X)(f \circ \varphi^t(x))$$

since

$$\frac{d}{dt}(f \circ \varphi^t)(x) = df(\varphi^t(x))X(\varphi^t(x)) = df(f^{-1}f \circ \varphi^t(x))X(f^{-1}f \circ \varphi^t(x)) = f_*X(f \circ \varphi^t(x))$$

Proposition 3.1. *Let H, K be two autonomous $C^{1,1}$ Hamiltonians of the symplectic manifold M and assume $\{H, K\}$ to be C^0 small. Then denoting by φ^t, ψ^s the Hamiltonian flows of H and K , the Hamiltonian isotopy $t \mapsto \varphi^t \psi^s \varphi^{-t} \psi^{-s}$ is generated by a C^0 small (time-dependent) Hamiltonian.*

Proof. Indeed we have, setting $u = \varphi^t \psi^s \varphi^{-t} \psi^{-s}(x)$:

$$\begin{aligned} \frac{d}{dt}(\varphi^t \psi^s \varphi^{-t} \psi^{-s})(x) &= \frac{d}{dt} \varphi^t(\psi^s \varphi^{-t} \psi^{-s}(x)) + d(\varphi^t \psi^s)(\varphi^{-t} \psi^{-s}(x)) \left(\frac{d}{dt} \varphi^{-t} \right)(\psi^{-s}(x)) = \\ &X_H(\varphi^t \psi^s \varphi^{-t} \psi^{-s}(x)) - d(\varphi^t \psi^s)(\varphi^{-t} \psi^{-s}(x)) X_H(\varphi^{-t} \psi^{-s}(x)) = \\ &X_H(u) - d(\varphi^t \psi^s)(\psi^{-s} \varphi^{-t}(u)) X_H(\psi^{-s} \varphi^{-t}(u)) = \\ &[X_H - (\varphi^t \psi^s)_* X_H](u), \end{aligned}$$

and since for each symplectic diffeomorphism, ρ , we have $\rho_*X_L = X_{L\rho^{-1}}$, the vector field $[X_H - (\varphi^t\psi^s)_*X_H]$ is Hamiltonian, with Hamiltonian function

$$L_s(t, x) = H(x) - H(\psi^{-s}\varphi^{-t}(x))$$

We may thus compute

$$\frac{\partial}{\partial s}L_s(t, x) = -\frac{d}{ds}H(\psi^{-s}\varphi^{-t}(x)) = dH(\psi^{-s}\varphi^{-t}(x)) \cdot X_K(\psi^{-s}\varphi^{-t}(x)) = \{H, K\}(\psi^{-s}\varphi^{-t}(x))$$

and since $H(\varphi^{-t}(x)) = H(x)$, we have $L_0 = 0$ and we may then estimate

$$\begin{aligned} |L_s(t, x)| &= |L_s(t, x) - L_0(t, x)| \leq \int_0^s \left| \frac{\partial}{\partial \sigma} L_\sigma(t, x) \right| d\sigma \leq \\ &\int_0^s |\{H, K\}(\psi^{-\sigma}\varphi^{-t}(x))| d\sigma \leq s \|\{H, K\}\|_{C^0} \end{aligned}$$

Thus if $|\{H, K\}|$ is C^0 small, for each s , the flow $t \mapsto \varphi^t\psi^s\varphi^{-t}\psi^{-s}$ is generated by a C^0 small Hamiltonian. \square

Remember that according to definition 1.1, two autonomous Hamiltonians H and K **C^0 -commute** if and only if there exist sequences of smooth Hamiltonians H_n, K_n such that, in the C^0 topology: H_n goes to H , K_n goes to K , and $\{H_n, K_n\}$ goes to zero.

Fortunately this definition does not conflict with the standard one according to theorem 1.2 that we now prove:

Proof of theorem 1.2. Remember that all Hamiltonians are compact supported. We shall explain in appendix B how to extend this to more general situations.

We also assume temporarily that H, K are of class $C^{1,1}$. Let now H_n, K_n be a sequence of compact supported $C^{1,1}$ Hamiltonians, such that their C^0 -limits satisfy:

- (a). $\lim_{n \rightarrow \infty} H_n = H, \lim_{n \rightarrow \infty} K_n = K$
- (b). $\lim_{n \rightarrow \infty} \{H_n, K_n\} = 0$

We wish to prove that if H, K are $C^{1,1}$ then $\{H, K\} = 0$. Let φ_n^t, ψ_n^s the flows of H_n, K_n , and set $\rho_n(s, t, x) = \varphi_n^t\psi_n^s\varphi_n^{-t}\psi_n^{-s}(x)$. The flow $t \mapsto \rho_n(s, t, x)$ is the flow of the Hamiltonian $L_s^n(t, x)$.

Making use of the topology of c -convergence from definition 2.4 above and applying Corollary 2.8 we have that φ_n^t c -converges to φ^t , and ψ_n^s to ψ^s . Using the triangle inequality for $\tilde{\gamma}$ we see that $\varphi_n^t\psi_n^s\varphi_n^{-t}\psi_n^{-s}$ c -converges to $\varphi^t\psi^s\varphi^{-t}\psi^{-s}$. On the other hand, according to proposition 3.1, since $\{H_n, K_n\}$ is C^0 small, $\tilde{\gamma}(\varphi_n^t\psi_n^s\varphi_n^{-t}\psi_n^{-s})$ goes to zero and thus, using proposition 2.6, we get that $\varphi^t\psi^s\varphi^{-t}\psi^{-s} = \text{Id}$. Since this holds for any s, t , it obviously implies that H and K commute. This concludes our proof. \square

The proof required the Hamiltonians to be $C^{1,1}$ in order to define the flows of X_H, X_K . One should be able to deal with the slightly more general C^1 case by using the methods of [Humilière].

The same proof shows that the following definition of commutation is also compatible with the standard one

Definition 3.2. Let H, K be two C^0 Hamiltonians on T^*N . We shall say that H and K **c -commute** if and only if there exists sequences H_n, K_n such that denoting the flows of H_n and H by φ_n^t, φ^t , and those of K_n and K by ψ_n^t, ψ^t (since H, K are only C^0 , the flows are only defined in $\mathfrak{H}\mathfrak{D}_\gamma(M)$), we have

$$\begin{aligned} \varphi_n^t &\xrightarrow{c} \varphi^t \\ \psi_n^t &\xrightarrow{c} \psi^t \\ \varphi_n^t \psi_n^s \varphi_n^{-t} \psi_n^{-s} &\xrightarrow{c} Id \end{aligned}$$

Note that even though this is a very natural notion, it is not the one we will need to generalize theorem 1.2.

3.1 The time-dependent and multi-time case

3.1.1 The time-dependent case

Assume now H, K are time-dependent, that is of the form $H(t, z)$ and $K(t, z)$. Commutation is then expressed by different formulas. If X_H, X_K are the Hamiltonian vector fields, we define

$$[H, K](t, z) = d_z H(t, z) X_K(t, z) + \frac{\partial H}{\partial t}(t, z) - \frac{\partial K}{\partial t}(t, z)$$

In the sequel, even for time-dependent vector fields we denote by $\{H, K\}(t, z)$ the Poisson bracket of H, K for fixed time, that is $\{H, K\}(t, z) = d_x H(t, z) X_K(t, z)$, so that

$$[H, K](t, z) = \{H, K\}(t, z) + \frac{\partial H}{\partial t}(t, z) - \frac{\partial K}{\partial t}(t, z)$$

The vanishing of $[H, K]$ has the following interpretation. To $H(t, z)$ we may associate the autonomous Hamiltonian

$$\tilde{H}(t, \tau, z) = \tau + H(t, z)$$

defined on $(\mathbb{R}^2 \times M, d\tau \wedge dt + \omega)$, and the associated flow is

$$\Phi^s : (t, \tau, z) \mapsto (t + s, \tau + H(t, z) - H(t + s, \varphi_t^{t+s}(z)), \varphi_t^{s+t}(z))$$

where φ_t^{t+s} is the flow on M of the non autonomous Hamiltonian $H(t, x)$.

Similarly to K we associate Ψ^s . Now it is easy to check that $[H, K](t, z) = \{\tilde{H}, \tilde{K}\}(t, \tau, z)$, the commutation of Φ^t and Ψ^s implies obviously that

$$\psi_b^d \varphi_a^b(z) = \varphi_c^d \psi_a^c(z)$$

for any a, b, c, d with $d - b = c - a$.

With this definition, the results of the previous section can be extended to the time-dependent case.

Definition 3.3. Let $H(t, z), K(t, z)$ be two continuous Hamiltonians. We shall say that H and K C^0 -**commute** if and only if there are sequences H_n, K_n such that $[H_n, K_n]$ goes to zero in the C^0 topology.

If H, K are of class $C^{1,1}$ this definition coincides with the condition $[H, K] = 0$.

3.1.2 The multi-time case

We shall limit ourselves to the case $d = 2$. We consider $H(s, t, z), K(s, t, z)$ and we consider the Hamiltonian flows

$$\frac{d}{dt} \varphi_{t_0}^t(s; x) = X_H(s, t, \varphi_{t_0}^t(s; x))$$

$$\frac{d}{ds} \psi_{s_0}^s(t; x) = X_K(s, t, \psi_{s_0}^s(t; x))$$

Then, we consider the Hamiltonians on $T^*(N \times \mathbb{R}^2)$ given by

$$\tilde{H}(s, t, \sigma, \tau, x) = \tau + H(s, t, z)$$

$$\tilde{K}(s, t, \sigma, \tau, x) = \sigma + K(s, t, z)$$

Denote their flows by $\tilde{\Phi}^a, \tilde{\Psi}^b$ respectively. The last component of $\tilde{\Phi}^a(s, t, \sigma, \tau)$ is given by $\varphi_t^{t+a}(s; z)$ and the last component of $\tilde{\Psi}^b(s, t, \sigma, \tau)$ by $\psi_s^{s+b}(t; z)$.

Now

$$\{\tilde{H}, \tilde{K}\} = \{H, K\}(s, t, z) + \frac{\partial H}{\partial t}(s, t, z) - \frac{\partial K}{\partial s}(s, t, z)$$

and the commutation of the flows $\varphi_{s_0}^s(t; x)$ and $\psi_{t_0}^t(s; x)$ is given by the vanishing of the above expression.

Definition 3.4. We set

$$\ll H, K \gg = \{H, K\}(s, t, z) + \frac{\partial H}{\partial t}(s, t, z) - \frac{\partial K}{\partial s}(s, t, z)$$

Proposition 3.5. The “flows” $\varphi_{s_0}^s(t; z)$ and $\psi_{t_0}^t(s; z)$ commute, or more explicitly

$$\varphi_{s_0}^s(t; \psi_{t_0}^t(s_0; z)) = \psi_{t_0}^t(s; \varphi_{s_0}^s(t_0; z))$$

if and only if $\ll H, K \gg = 0$.

Remark 3.6. By applying Proposition 3.1, we get that all our different notions of commutation (in the time-dependent or multi-time case) are well defined in the continuous case, and the definition is compatible with the usual definition in the $C^{1,1}$ case.

4 Multi-time Hamilton-Jacobi equations and the proof of theorem 1.3

Consider the system of equations (MHJ) defined in the introduction. For small t_1, \dots, t_d , and a Hamiltonian of class $C^{1,1}$, we can find a solution to the above equation provided the Hamiltonians commute in the sense of the previous section, that is the functions $\tau_j + H_j(t_1, \dots, t_d, x, p)$ commute on $T^*(\mathbb{R}^d \times N)$, which we can rewrite as:

$$(H1) \quad \{H_j, H_k\}(t, z) + \frac{\partial H_j}{\partial t_k}(t, z) - \frac{\partial H_k}{\partial t_j}(t, z) = 0 \quad t \in \mathbb{R}^d, z \in T^*N$$

4.1 The smooth case

Under assumption (H1) it is easy to check that the method of characteristics will yield a solution for t_j small.

Let us remind the reader of the geometric approach to this method. Set

$$\Lambda_{f; H_1, \dots, H_d} = \{(0, \dots, 0, x, -H_1(0, \dots, 0, x, df(x)), \dots, -H_d(0, \dots, 0, x, df(x)), df(x)) \mid x \in N\}$$

This is an isotropic submanifold in $T^*(N \times \mathbb{R}^d)$, and the vector fields X_j associated to $\tilde{H}_j(t_1, \dots, t_d, \tau_1, \dots, \tau_d, x, p) = \tau_j + H_j(t_1, \dots, t_d, x, p)$ being pairwise in involution, are all tangents to the submanifold

$$\Sigma = \bigcap_{j=1}^d \tilde{H}_j^{-1}(0)$$

and transverse to $\Lambda_{f;H_1,\dots,H_d}$. As a result the flows $\tilde{\Phi}_j^t$ of X_j commute and defining

$$L_{f;H_1,\dots,H_d} = \bigcup_{t_1,\dots,t_d \in \mathbb{R}^d} \tilde{\Phi}_1^{t_1} \dots \tilde{\Phi}_j^{t_j} \dots \tilde{\Phi}_d^{t_d}(\Lambda_{f;H_1,\dots,H_d})$$

the commutation property insures that the order in which we take the $\tilde{\Phi}_j^{t_j}$ is irrelevant, and that $L_{f;H_1,\dots,H_d}$ is a Lagrangian submanifold contained in $H_j^{-1}(0)$, hence in Σ .

The following lemma is a straightforward extension of a result from [Viterbo-Ottolenghi].

Lemma 4.1. *The manifold $L_{f;H_1,\dots,H_d}$ is Hamiltonianly isotopic to the zero section in $T^*(\mathbb{R}^d \times N)$.*

Proof. Consider the family $\Phi_{j,\lambda}^\tau$ of flows of the Hamiltonian $\tau_j + \lambda H(t_1, \dots, t_d, x, p)$. We have $\Phi_{j,1}^\tau = \Phi_j^\tau$ and $\Phi_{j,0}^\tau$ is the translation by τ in the t_j coordinate.

Now the family $\Phi_{j,\lambda}^\tau$ is a continuous family of proper maps, and the family of Lagrangian submanifolds given by

$$L_{f;H_1,\dots,H_d}^\lambda = \bigcup_{t_1,\dots,t_d \in \mathbb{R}^d} \tilde{\Phi}_{1,\lambda}^{t_1} \dots \tilde{\Phi}_{j,\lambda}^{t_j} \dots \tilde{\Phi}_{d,\lambda}^{t_d}(\Lambda_{\lambda f;H_1,\dots,H_d})$$

is an exact proper Hamiltonian isotopy of Lagrangian submanifolds.

Note that we do not need here that the Hamiltonians commute, and we shall indeed use the lemma in the more general setting. \square

Now, still assuming the H_j are of class $C^{1,1}$ in (x, p) , section 2 implies that $L_{f;H_1,\dots,H_d}$ has a GFQI. We thus have a variational solution $u_{f;H_1,\dots,H_d}$ of the following equations

$$(\star) \begin{cases} \frac{\partial}{\partial t_1} u(t_1, t_2, \dots, t_d, x) + H_1(t_1, t_2, \dots, t_d, x, \frac{\partial}{\partial x} u(t_1, t_2, \dots, t_{d-1}, t_d, x)) = 0 \\ \frac{\partial}{\partial t_2} u(0, t_2, \dots, t_d, x) + H_2(0, t_2, \dots, t_d, x, \frac{\partial}{\partial x} u(0, t_2, \dots, t_{d-1}, t_d, x)) = 0 \\ \vdots \\ \frac{\partial}{\partial t_d} u(0, 0, \dots, t_d, x) + H_d(0, 0, \dots, t_d, x, \frac{\partial}{\partial x} u(0, 0, \dots, 0, t_d, x)) = 0 \end{cases}$$

The commutation of the $\tilde{\Phi}_j^{t_j}$ implies that for any permutation σ of $\{1, \dots, n\}$, the Lagrangian manifold

$$L_{f;H_1,\dots,H_d}^\sigma = \bigcup_{(t_1,\dots,t_q) \in \mathbb{R}^q} \tilde{\Phi}_{\sigma(1)}^{t_{\sigma(1)}} \dots \tilde{\Phi}_{\sigma(q)}^{t_{\sigma(q)}}(\Lambda_{f;H_1,\dots,H_d})$$

coincides with

$$L_{f;H_1,\dots,H_d}$$

Thus the associated function $u_{f;H_1,\dots,H_d}^\sigma = u_{f;H_{\sigma(1)},\dots,H_{\sigma(d)}}$ coincides with $u_{f;H_1,\dots,H_d}$. As a result, u is a solution of any equation obtained by replacing $\frac{\partial}{\partial t_j}$ by $\frac{\partial}{\partial t_{\sigma(j)}}$ and H_j by $H_{\sigma(j)}$. Looking at the first equation, we see that

$$\frac{\partial}{\partial t_{\sigma(1)}}u(t_1, t_2, \dots, t_d, x) + H_{\sigma(1)}(t_1, t_2, \dots, t_d, x, \frac{\partial}{\partial x}u(t_1, t_2, \dots, t_{d-1}, t_d, x)) = 0$$

and since $\sigma(1)$ is arbitrary, we see that u solves (MHJ).

Our problem is thus solved for initial data and equation of class $C^{1,1}$.

However we are really interested in the C^0 case. This is the subject of the next section.

4.2 The continuous case

To deal with the C^0 case, we must first extend the construction of the Lagrangian submanifold

$$L_{f;H_1,\dots,H_d}$$

to general Hamiltonians and initial conditions. Note that since the Hamiltonians do not commute anymore, order now matters. We shall consider for each permutation σ of $\{1, \dots, d\}$, the manifold

$$L_{f;H_1,\dots,H_d}^\sigma$$

and as in Lemma 4.1, the manifold $L_{f;H_1,\dots,H_d}^\sigma$ is Hamiltonianly isotopic to the zero section, thus to each such Lagrangian we may associate a unique function u^σ .

If the H_j are only C^0 and we consider a sequence H_j^ν of C^2 Hamiltonians converging to H_j . We shall then associate to these a sequence of Lagrangian submanifolds and a sequence of functions u_ν^σ , which for $\sigma = Id$, is the solution of the set of equations obtained by replacing in (\star) the H_j by H_j^ν .

We thus want to prove that the conditions $\lim_{\nu \rightarrow \infty} H_j^\nu = H_j$, and $\lim_{\nu \rightarrow \infty} \{H_j^\nu, H_k^\nu\} = 0$, imply

$$\lim_{\nu \rightarrow \infty} |u_\nu^\sigma - u_\tau^\nu| = 0$$

for any permutations σ, τ . One should notice that this does not say much about the $L_{f;H_1^\nu,\dots,H_d^\nu}^\sigma$ and in particular we do not claim their convergence in the C^0 sense.

Proposition 4.2. *Assume $\lim_{\nu \rightarrow \infty} f^\nu = f$, $\lim_{\nu \rightarrow \infty} H_j^\nu = H_j$ and $\lim_{\nu \rightarrow \infty} \{H_j^\nu, H_k^\nu\} = 0$. Then $u_{f;H_1,\dots,H_d} = \lim_{\nu \rightarrow \infty} u_\nu^\sigma$ is independent of σ and is a solution of the multi-time Hamilton-Jacobi equation (MHJ). Moreover $u_{f;H_1,\dots,H_d}$ is a variational solution of each single equation. If the H_j are convex in p , then according to [Zhukovskaya] variational and viscosity solutions do coincide and thus $u_{f;H_1,\dots,H_d}$ is the viscosity solution of each single equation.*

The proof of this proposition will follow from a more general theorem, involving c -convergence. From now on, we shall only consider the case $d = 2$, since the general case is in all respects similar.

4.3 Capacity for manifolds with boundary

In the sequel we shall need to define γ on \mathcal{L} in the symplectic manifold $T^*([0, 1] \times N)$ and $\mathcal{D}_\omega(T^*([0, 1]^2 \times N))$ or more generally for the case of the cotangent bundle of a manifold M with boundary. Let M be a manifold with boundary, and L be a Lagrangian submanifold in T^*N such that $\partial L = L \cap T_{\partial N}^*N$. Assuming moreover the intersection to be transverse, we can consider the manifold $\tilde{L} \subset T^*(\tilde{N})$ where \tilde{L}, \tilde{N} are the doubles of L, N . We shall write $\tilde{N} = N_- \cup N_+, \tilde{L} = L_- \cup L_+$ and we identify L, N with L_-, N_- . We also denote by τ be the involution of \tilde{N} sending N_- to N_+ , and by t the coordinate defining the collar of ∂M . By assumption, the reduction of L by $\{t = t_0\}$, denoted by L_{t_0} is a Lagrangian regular homotopy. We may modify this family near $\{t = 0\}$, so that it is constant for t close to 0. Then near $\{t = 0\}$, L is uniquely defined by the real function $\tau(t, z)$ defined on $[0, \varepsilon] \times L_0$ such that $L = \{(t, \tau(t, z), z) \mid z \in L_t\}$. Then $\tau(t, z) = \tau(z)$ near $t = 0$, and we may thus extend τ and L_t to $[-\varepsilon, \varepsilon] \times L_0$, so that $\tau(-t, z) = \tau(t, z)$ and $L_{-t} = L_t$. This way we may extend L to a Lagrangian manifold $\tilde{L} \subset T^*\tilde{M}$, uniquely defined by L .

Notice that if L is Hamiltonianly isotopic to the zero section, by a Hamiltonian preserving $T_{\partial N}^*N$, then clearly the same holds for \tilde{L} . Therefore \tilde{L} has a GFQI, and we can compute $\gamma(\tilde{L})$. We define $\gamma(L) = \gamma(\tilde{L})$. Note also that if $S(x, \xi)$ is a GFQI for \tilde{L} , then $(\tau^*S)(x, \xi) = S(\tau(x), \xi)$ also generates \tilde{L} . In fact any S generating L over T^*N has such an extension.

4.4 S -commuting Hamiltonians

Remember that we denoted by $\mathcal{H}(M)$ be the set of smooth time-dependent Hamiltonians on $\mathbb{R} \times M$, and by $\mathcal{H}\mathcal{D}(M)$ the set of time one maps of such Hamiltonians, and defined $\mathfrak{H}_\gamma(M)$, the completion of $\mathcal{H}(M)$ for the metric γ , and $\mathfrak{H}\mathcal{D}_\gamma(M)$ the completion of $\mathcal{H}\mathcal{D}(M)$ for the metric γ .

As we pointed out, the ‘‘time one flow’’ map $\mathcal{H}(M) \xrightarrow{\sim} \mathcal{H}\mathcal{D}(M)$ extends to a continuous map $\mathfrak{H}_\gamma(M) \rightarrow \mathfrak{H}\mathcal{D}_\gamma(M)$, and in particular is well defined on the set of continuous maps $C^0(\mathbb{R} \times M)$.

Let φ^s, ψ^t be the flows of the autonomous Hamiltonians H and K . Then we introduce the following commutation condition. Consider $\rho^{(s,t)} = \varphi^s \psi^t \varphi^{-s} \psi^{-t}$ for $(s, t) \in [0, 1]^2$, and the map

$$C(H, K) : (s, t, \sigma, \tau, z) \rightarrow (s, t, \tilde{\sigma}(s, t, z), \tilde{\tau}(s, t, z), \rho^{(s,t)}(z))$$

from $T^*([0, 1]^2 \times N)$ into itself. We may choose $\tilde{\sigma}, \tilde{\tau}$ such that $C(H, K)$ is a Hamiltonian diffeomorphism, using the fact that $\lambda \rightarrow \rho^{(s, \lambda t)}$ is an Hamiltonian isotopy starting from the identity. The functions $\tilde{\sigma}, \tilde{\tau}$ are well defined up to the addition of $\frac{\partial c}{\partial s}(s, t)$ and $\frac{\partial c}{\partial t}(s, t)$ respectively, where c is some C^1 function.

If we are given Hamiltonians $H(z), K(z)$ generating φ^s, ψ^t , we can assume $s \rightarrow \rho^{(s, t)}$ is generated by

$$L_t(s, z) = H(z) - H(\psi^{-t}\varphi^{-s}(z))$$

We may also compute

$$\frac{\partial}{\partial t} L_t(s, z) = \{H, K\}(\psi^{-t}\varphi^{-s}(z))$$

and note that

$$\tilde{\tau}_1(t_1, t_2, z) = \tau_1 - L_{t_2}(t_1, \varphi_1^{t_1}\varphi_2^{t_2}\varphi_1^{-t_1}\varphi_2^{-t_2}(z)) = H_1(\varphi_1^{t_1}\varphi_2^{t_2}\varphi_1^{-t_1}\varphi_2^{-t_2}(z)) - H_1(\varphi_1^{-t_1}\varphi_2^{-t_2}(z))$$

$$\tilde{\tau}_2(t_1, t_2, z) = \tau_2 - t_1 \int_0^1 \{H_1, H_2\}(\varphi_1^{-\sigma t_1}\varphi_2^{-t_2}(z)) d\sigma$$

Note that if either M is compact and we impose the condition $\int_M H_j(z)\omega^n$ or H_j are compact supported, then $L_{t_2}(t_1, z)$ will satisfy the same condition, and this defines uniquely $C(H, K)$ as the time one map of a Hamiltonian satisfying this normalization condition.

Definition 4.3. (a). We shall say that φ^s, ψ^t S -commute up to ε , if and only if $\gamma(C(H, K)) \leq \varepsilon$.

(b). Let H, K be in $\mathfrak{H}\mathfrak{D}_\gamma(T^*N)$. We say that they S -commute, if and only if there are sequences H_j, K_j such that H_j, K_j converge to H, K (for the metric γ) and $\gamma(C(H_j, K_j))$ goes to zero.

We now have

Lemma 4.4. (a). If φ_1^t, φ_2^t S -commute up to ε then $\gamma(\rho^{(t_1, t_2)}) \leq \varepsilon$ for all pairs $(t_1, t_2) \in [0, 1]^2$. In particular S -commuting Hamiltonians c -commute.

(b). If for all (t_1, t_2) in $[0, 1]^2$, the Hofer norm of $\rho^{(t_1, t_2)}$, is less than ε , then φ_1^t, φ_2^t c -commute up to ε . In particular C^0 -commuting Hamiltonians S -commute.

Proof. The first statement follows easily by symplectic and the reduction inequality (see [Viterbo 1], page 705, prop. 5.1) that can be stated as follows:

Let $S(x, y, \xi)$ be a GFQI, and denote by S_x the function $S_x(y, \xi) = S(x, y, \xi)$.

Then

$$c(1, S) \leq \inf_x c(1_x, S_x) \leq \sup_x c(\mu_x, S) \leq c(\mu, S)$$

The second statement follows from the proof of the inequality

$$d(U) \leq 2 \sup_x d(U_x)$$

in [Viterbo 3], where d is the displacement energy, i.e.

$$d(U) = \inf\{|H|_{C^0} \mid \varphi_H^1(U) \cap U = \emptyset\}$$

□

4.5 Different suspensions of Hamiltonian isotopies

Let $H(t, z)$ be a time-dependent Hamiltonian on T^*N . We may associate to H the Hamiltonians on $T^*(\mathbb{R} \times N)$ given by $\tilde{H}(t, \tau, z) = \tau + H(t, z)$ and $\hat{H}(s; t, \tau, z) = sH(st, z)$, with respective flows

$$\Phi^s : (t, \tau, z) \rightarrow (t + s, \tau - H(t, \varphi_t^{t+s}(z)), \varphi_t^{t+s}(z))$$

$$\hat{\Phi}^s : (t, \tau, z) \rightarrow (t, \tau - sH(st, \varphi_0^{st}(z)), \varphi_0^{st}(z))$$

Note that if H is C^0 small, we have, denoting by T_s the translation on the t variable, $\gamma(\Phi^s T_{-s})$ and $\gamma(\hat{\Phi}^s)$ are small, in other words, Φ^s is c -close to T_s and $\hat{\Phi}^s$ is c -close to Id .

5 Proof of theorem 1.3.

The construction of variational solutions for a single equation can be rephrased as follows. Let Λ_0 be a Lagrangian submanifold of T^*N , and $H(t, z)$ be a time-dependent Hamiltonian. We consider $\tilde{\Lambda}_{0,H} = \{(0, -H(0, z), z) \mid z \in \Lambda_0\}$, and $L = \bigcup_{t \in \mathbb{R}} \Phi^t(\Lambda_{0,H})$.

We can obtain L as follows. Consider the map

$$\begin{aligned} F : T^*(\mathbb{R} \times \mathbb{R} \times N) &\rightarrow T^*(\mathbb{R} \times \mathbb{R} \times N) \\ (s, \sigma, \zeta) &\mapsto (s, \tilde{\sigma}, \Phi^s(\zeta)) \end{aligned}$$

and $L_{0,H}$ any Lagrangian submanifold in $T^*(\mathbb{R} \times N)$ such that $\tilde{\Lambda}_{0,H} \subset L_{0,H}$. Let $t : T^*(\mathbb{R} \times N) \rightarrow \mathbb{R}$ be the projection on the first coordinate of $\mathbb{R} \times N$.

Proposition 5.1. L is the reduction of $F(L_{0,H})$ by the coisotropic submanifold $\{s = t\}$.

Proof. Indeed, we look for the set of $(s, \tilde{\sigma}, \Phi^s(0, -H(0, z), z))$ such that $t = s$. But $\Phi^s(t, \tau, z) = (t + s, \tau + H(t, z) - H(t + s, \varphi_t^{t+s}(z)), \varphi_t^{t+s}(z))$. Intersecting with $t = s$ means that we consider the points $\Phi^s(0, \tau, z)$ where $(0, \tau, z) \in L_{0,H}$ and by assumption this means that $(0, \tau, z) \in \tilde{\Lambda}_{0,H}$. \square

Now we will first estimate $\gamma(L_{0,H})$ and then deduce the inequality

$$\gamma(L) \leq \tilde{\gamma}(F) + \gamma(L_{0,H})$$

Lemma 5.2. Any isotopy Ξ^s of $\{t = 0\}$ in $T^*(\mathbb{R} \times N)$ can be generated by an arbitrarily small Hamiltonian preserving the hypersurfaces $\{t = t_0\}$.

Proof. Indeed if we have an isotopy of the type $(0, \tau, x, p) \mapsto (0, \tau + \tau^s(x, p), x, p)$, it is induced by a Hamiltonian $H(t, x, p)$ and only depends on $\frac{\partial H}{\partial t}(0, x, p) = \frac{\partial}{\partial s} \tau^s(x, p)$. The extension of H to $T^*(\mathbb{R} \times N)$ can be arbitrarily small, for example we may take $H(s; t, \tau, x, p) = \chi(t) \frac{\partial}{\partial s} \tau^s(x, p)$, where $\dot{\chi}(0) = 1$ and χ has arbitrarily small C^0 norm. \square

Proposition 5.3. For any positive ε , we may choose $L_{0,H}$ so that we have

$$\gamma(L_{0,H}) \leq \gamma(\Lambda_0) + \varepsilon$$

Remark 5.4. Note that necessarily we have $\gamma(L_{0,H}) \geq \gamma(\Lambda_0)$. This follows from the reduction inequality.

Proof. We notice that $L_{0,0} = \mathbb{R} \times \{0\} \times \Lambda_0$ satisfies obviously $\gamma(L_{0,0}) = \gamma(\Lambda_0)$. Using the notation $L_{0,0} \cap \{t = 0\} = \tilde{\Lambda}_{0,0}$, the previous lemma allows us to find a Hamiltonian isotopy, Ξ^s on $T^*(\mathbb{R} \times N)$ such that

- Ξ^s preserves $t = 0$
- on $t = 0$, we have $\Xi^1(\tilde{\Lambda}_{0,0}) = \tilde{\Lambda}_{0,H}$
- Ξ^s is generated by an arbitrarily C^0 small Hamiltonian.

Therefore

$$\gamma(L_{0,H}) = \gamma(\Xi^1(\Lambda_{0,0})) \leq \tilde{\gamma}(\Xi^1) + \gamma(\Lambda_{0,0}) \leq \varepsilon + \gamma(\Lambda_{0,0})$$

\square

Corollary 5.5. *We have $\gamma(L) \leq \tilde{\gamma}(F) + \gamma(\Lambda_0)$. Moreover if L_1 is the Lagrangian submanifold associated to H_1, Λ_1 and L_2 to H_2, Λ_2 , we have*

$$\gamma(L_1 - L_2) \leq \tilde{\gamma}(F_1 \circ F_2^{-1}) + \gamma(\Lambda_1 - \Lambda_2)$$

Proof. The first inequality follows from the second by setting

$$L_1 = L, F_1 = F, \Lambda_1 = \Lambda, L_2 = 0_{\mathbb{R} \times N}, F_2 = \text{Id}, \Lambda_2 = O_N$$

Since L_1 and L_2 are the reductions of $F_1(L_{1,H_1})$ and $F_2(L_{1,H_1})$, using the reduction inequality it will be enough to prove that

$$\gamma(F_1(L_{1,H_1}) - F_2(L_{2,H_2})) \leq \tilde{\gamma}(F_1 \circ F_2^{-1}) + \gamma(\Lambda_1 - \Lambda_2)$$

Since the map $F_2 \circ F_1^{-1}$ sends $F_1(L_{1,H_1})$ to $F_2(L_{1,H_1})$, we have by the triangle inequality and the symplectic invariance of γ ,

$$\begin{aligned} \gamma(F_1(L_{1,H_1}) - F_2(L_{2,H_2})) &\leq \gamma(F_1(L_{1,H_1}) - F_2(L_{1,H_1})) + \gamma(F_2(L_{1,H_1}) - F_2(L_{2,H_2})) \leq \\ &\tilde{\gamma}(F_2 \circ F_1^{-1}) + \gamma(L_{1,H_1} - L_{2,H_2}) \end{aligned}$$

so that using invariance by reduction, we have

$$\gamma(L_1 - L_2) \leq \tilde{\gamma}(F_2 \circ F_1^{-1}) + \gamma(L_{1,H_1} - L_{2,H_2})$$

Finally we claim that for any positive ε we may find L_{1,H_1}, L_{2,H_2} such that $\gamma(L_{1,H_1} - L_{2,H_2}) \leq \gamma(\Lambda_1 - \Lambda_2) + \varepsilon$. Indeed remember that L_{1,H_1}, L_{2,H_2} are not uniquely defined, but just required to be Lagrangian submanifolds contained in $\tau + H_1(t, z) = 0$ and $\tau + H_2(t, z) = 0$ respectively, with reductions L_1, L_2 . Since $\gamma(L_{1,0} - L_{2,0}) = \gamma(\Lambda_1 - \Lambda_2)$ and we saw in Lemma 5.2 that we may go from $L_{1,0}$ to L_{1,H_1} by applying a Hamiltonian with C^0 norm less than ε , and similarly from $L_{2,0}$ to L_{2,H_1} we get that $\gamma(L_{1,H_1} - L_{1,0})$ and $\gamma(L_{2,H_2} - L_{2,0})$ are arbitrarily small. We may thus conclude

$$\gamma(L_{1,H_1} - L_{2,H_2}) \leq \gamma(L_{1,H_1} - L_{1,0}) + \gamma(L_{1,0} - L_{2,0}) + \gamma(L_{2,0} - L_{2,H_2}) \leq 2\varepsilon + \gamma(\Lambda_1 - \Lambda_2)$$

Since ε is arbitrarily small, we get the announced inequality. \square

Our goal here is to prove the following theorem

Theorem 5.6. *Assume the Hamiltonians $H_1(t_1, \dots, t_d, x, p), \dots, H_d(t_1, \dots, t_d, x, p)$ are Lipschitz, and S -commute. Then equation (MHJ) has a unique variational solution. If the H_j are convex in p , then u is a viscosity solution of each individual equation.*

We limit ourselves to the case $d = 2$.

We consider the flows Φ_1^s, Φ_2^s associated to $\tau_1 + H_1(t_1, t_2, z), \tau_2 + H_2(t_1, t_2, z)$, and associated isotopies depending on 2 parameters. More precisely $\varphi_1^{(t_1, t_1+s)}(t_2; z)$ is the flow of $(t_1, z) \rightarrow H(t_1, t_2, z)$ between times t_1 and $t_1 + s$ (t_2 being considered as fixed). Similarly $\varphi_2^{(t_2, t_2+s)}(t_1; z)$ is the flow associated to the Hamiltonian $(t_2, z) \rightarrow H_2(t_1, t_2, z)$ between times t_2 and $t_2 + s$.

Then Φ_1^s, Φ_2^s are symplectomorphisms from $T^*(\mathbb{R}^2 \times N)$ given by

$$\Phi_1^s(t_1, t_2, \tau_1, \tau_2, z) = (t_1 + s, t_2, \tau_1 + \hat{\tau}_1^s(t_1, t_2, z), \tau_2 + \hat{\tau}_2^s(t_1, t_2, z), \varphi_1^{(t_1, t_1+s)}(t_2; z))$$

$$\Phi_2^s(t_1, t_2, \tau_1, \tau_2, z) = (t_1 + s, t_2, \tau_1 + \check{\tau}_1^s(t_1, t_2, z), \tau_2 + \check{\tau}_2^s(t_1, t_2, z), \varphi_2^{(t_2, t_2+s)}(t_1; z))$$

Note that since $L_1 = \bigcup_{t_1 \in \mathbb{R}} \Phi_1^{t_1}(\tilde{\Lambda}_0)$ is not contained in the hypersurface $\tau_2 + H_2(t_1, t_2, x, p) = 0$, in order to get a Lagrangian submanifold after applying the flow Φ_2^s we must move L_1 in the $\frac{\partial}{\partial \tau_2}$ direction, so that its image is included in this hypersurface.

Setting

$$\tau_2^s(z) = - \int_0^s \frac{\partial H_1}{\partial t_2}(\sigma, 0, \varphi_1^{(0, \sigma)}(0; z)) d\sigma$$

and

$$\tilde{L}_1 = \{(t_1, 0, -H_1(t_1, 0, \varphi^{(0, t_1)}(z)), -H_2(0, 0, z) + \tau_2^{t_1}(z), \varphi_1^{(0, t_1)}(0; z)) \mid z = (x, df(x)) \in T^*N\}$$

we may indeed check that on $\tilde{L}_1, \tau_2 + H_2(t_1, t_2, z)$ vanishes (note that $\tilde{\Lambda}_1$ is contained in $\{t_2 = 0\}$).

We thus define a “normalization map”, defined over $\{t_2 = 0\}$:

$$V_{1,2} : (t_1, 0, \tau_1, \tau_2, z) \mapsto (t_1, 0, \tau_1, -H_2(t_1, 0, z), z)$$

and similarly

$$V_{2,1} : (0, t_2, \tau_1, \tau_2, z) \mapsto (0, t_2, -H_1(t_1, 0, z), \tau_2, z)$$

These maps are not diffeomorphisms, but their restriction to \tilde{L}_1 (resp. \tilde{L}_2) is one, and since it preserves the Liouville form $(\tau_2 dt_2 + \tau_1 dt_1 + \lambda)$, its restriction to \tilde{L}_1 (resp. \tilde{L}_2) extends to a Hamiltonian isotopy $\tilde{V}_{1,2}$ (resp. $\tilde{V}_{2,1}$) which according to lemma 5.2 are generated by arbitrarily C^0 -small Hamiltonians ⁸. We will use the maps

⁸Note that $\tilde{V}_{1,2}$ and $\tilde{V}_{2,1}$ depend on \tilde{L}_1 and \tilde{L}_2 respectively.

$$W_{1,2} : (s_1, s_2, \sigma_1, \sigma_2, \zeta) \mapsto (s_1, s_2, \hat{\sigma}_1, \hat{\sigma}_2, \Phi_2^{s_2} \tilde{V}_{1,2} \Phi^{-s_2})$$

and

$$W_{2,1} : (s_1, s_2, \sigma_1, \sigma_2, \zeta) \mapsto (s_1, s_2, \check{\sigma}_1, \check{\sigma}_2, \Phi_1^{s_1} \tilde{V}_{2,1} \Phi_1^{-s_1})$$

also generated by C^0 -small Hamiltonians.

Now we claim that the following submanifolds are Lagrangian

$$L_{1,2} = \bigcup_{(s_1, s_2) \in \mathbb{R}^2} \Phi_1^{s_1} \tilde{V}_{2,1} \Phi_2^{s_2} (\Lambda_0)$$

$$L_{2,1} = \bigcup_{(s_1, s_2) \in \mathbb{R}^2} \Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} (\Lambda_0)$$

It will later be useful to notice that

$$\Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} \Phi_2^{-s_2} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1} = \left(\Phi_2^{s_2} \tilde{V}_{1,2} \Phi_2^{-s_2} \right) \circ \left(\Phi_2^{s_2} \Phi_1^{s_1} \Phi_2^{-s_2} \Phi_1^{-s_1} \right) \circ \left(\Phi_1^{s_1} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1} \right)$$

We then consider the map C as above:

$$\begin{aligned} T^*(\mathbb{R}^2 \times \mathbb{R}^2 \times N) &\longrightarrow T^*(\mathbb{R}^2 \times \mathbb{R}^2 \times N) \\ (s_1, s_2, \sigma_1, \sigma_2, \zeta) &\longrightarrow (s_1, s_2, \tilde{\sigma}_1, \tilde{\sigma}_2, \Phi_1^{s_1} \Phi_2^{s_2} \Phi_1^{-s_1} \Phi_2^{-s_2} (\zeta)) \end{aligned}$$

where $\zeta = (t_1, t_2, \tau_1, \tau_2, z)$, and the map $\tilde{C} = W_{1,2} \circ C \circ W_{2,1}^{-1}$.

We do not claim that \tilde{C} sends $L_{1,2}$ to $L_{2,1}$, since these manifolds are in $T^*(\mathbb{R}^2 \times N)$ and not $T^*(\mathbb{R}^2 \times \mathbb{R}^2 \times N)$.

However, let Δ be the coisotropic submanifold of $T^*(\mathbb{R}^2 \times \mathbb{R}^2 \times N)$ given by

$$\Delta = \{s_1 = t_1, s_2 = t_2\}$$

Proposition 5.7.

$$[\tilde{C}(0_{\mathbb{R}^2} \times L_{2,1})]_{\Delta} = L_{1,2}$$

Proof.

$$(0_{\mathbb{R}^2} \times L_{1,2}) = \{(s_1, s_2, 0, 0, \Phi_1^{t_1} \tilde{V}_{2,1} \Phi_2^{t_2} (\zeta)) \mid \zeta \in \tilde{\Lambda}_0\}$$

then

$$\begin{aligned} \tilde{C}(0_{\mathbb{R}^2} \times L_{2,1}) &= \{(s_1, s_2, \tilde{\sigma}_1, \tilde{\sigma}_2, \Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} \Phi_2^{-s_2} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1} (\zeta)) \mid \zeta \in L_{2,1}\} = \\ &= \{(s_1, s_2, \tilde{\sigma}_1, \tilde{\sigma}_2, (\Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} \Phi_2^{-s_2} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1}) \circ (\Phi_1^{t_1} \tilde{V}_{2,1} \Phi_2^{t_2}) (\zeta)) \mid \zeta \in \tilde{\Lambda}_0, (t_1, t_2) \in \mathbb{R}^2\} \end{aligned}$$

Now if $t_1(\zeta) = t_2(\zeta) = 0$ we have

$$t_1((\Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} \Phi_2^{-s_2} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1}) \circ (\Phi_1^{t_1} \tilde{V}_{2,1} \Phi_2^{t_2})(\zeta)) = t_1$$

$$t_2((\Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} \Phi_2^{-s_2} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1}) \circ (\Phi_1^{t_1} \tilde{V}_{2,1} \Phi_2^{t_2})(\zeta)) = t_2$$

Thus intersection of this with $\{s_1 = t_1, s_2 = t_2\}$ will be given by

$$\begin{aligned} & \{(s_1, s_2, \tilde{\sigma}_1, \tilde{\sigma}_2, \Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1} \Phi_2^{-s_2} \tilde{V}_{2,1}^{-1} \Phi_1^{-s_1} \circ \Phi_1^{s_1} \tilde{V}_{2,1} \Phi_2^{s_2}(\zeta) \mid \zeta \in \tilde{\Lambda}_0\} = \\ & \{(s_1, s_2, \tilde{\sigma}_1, \tilde{\sigma}_2, \Phi_2^{s_2} \tilde{V}_{1,2} \Phi_1^{s_1}(\zeta) \mid \zeta \in \tilde{\Lambda}_0\} \end{aligned}$$

and we must figure out its projection according to the equivalence relation given by

$$(s_1, s_2, \sigma_1, \sigma_2, t_1, t_2, \tau_1, \tau_2, z) \sim (s'_1, s'_2, \sigma'_1, \sigma'_2, t'_1, t'_2, \tau'_1, \tau'_2, z')$$

if and only if

$$s'_1 = s_1 = t_1 = t'_1, s_2 = s'_2 = t_2 = t'_2, \sigma_1 + \tau_1 = \sigma'_1 + \tau'_1, \sigma_2 + \tau_2 = \sigma'_2 + \tau'_2$$

Since the map

$$(s_1, s_2, \sigma_1, \sigma_2, t_1, t_2, \tau_1, \tau_2, z) \longrightarrow (s_1 - t_1, s_2 - t_2, \sigma_1, \sigma_2, t_1, t_2, \sigma_1 + \tau_1, \sigma_2 + \tau_2, z)$$

is a symplectomorphism, the reduction of $X = \tilde{C}(0_{\mathbb{R}^2} \times L_{2,1})$ is given by

$$X_{\Delta} = \{(s_1, s_2, \sigma_1 + \tau_1, \sigma_2 + \tau_2, z) \mid (s_1, s_2, \sigma_1, \sigma_2, s_1, s_2, \tau_1, \tau_2, z) \in X\}$$

and this yields $L_{1,2}$ as announced. \square

Proposition 5.8. *Let φ_1^t, φ_2^t S -commute up to ε , then we have*

$$\gamma(L_{1,2} - L_{2,1}) \leq \varepsilon$$

Proof. Indeed it is enough to prove that $\tilde{\gamma}(\tilde{C})$ is small by assumption (Φ^t, Ψ^s S -commute up to ε). Since $\gamma(C)$ is small, and $W_{1,2}, W_{2,1}^{-1}$ are generated by C^0 small Hamiltonians, $\gamma(W_{1,2}), \gamma(W_{2,1})$ are also small, and using the triangle inequality, we get that $\tilde{\gamma}(\tilde{C})$ is small.

Now if $\tilde{\gamma}(\tilde{C}) \leq \varepsilon$

$$\gamma(\tilde{C}(0_{\mathbb{R}^2} \times L_{2,1}) - 0_{\mathbb{R}^2} \times L_{1,2}) \leq \varepsilon$$

Using the reduction inequality we obtain

$$\gamma(\widetilde{C}(0_{\mathbb{R}^2} \times L_{2,1}) - 0_{\mathbb{R}^2} \times L_{1,2})_{\Delta} \leq \varepsilon$$

and according to the above proposition 5.7, we get

$$\gamma(L_{2,1} - L_{1,2}) \leq \varepsilon$$

□

Proof of theorem 1.3. Indeed we just proved that if φ_1^t, φ_2^t S -commute up to ε on $[0, T]$, we have $\gamma(L_{1,2} - L_{2,1}) \leq \varepsilon$ and thus, by the reduction inequality we have

$$|u_{1,2}(t, x) - u_{2,1}(t, x)| \leq \varepsilon$$

on $[0, T] \times N$.

Now assume we have sequences H_1^ν, H_2^ν such that H_1^ν c -converges to H_1 , H_2^ν c -converges H_2 , and H_1^ν, H_2^ν S -commute up to ε_ν where $\lim_{\nu \rightarrow 0} \varepsilon_\nu = 0$. Then we have

$$|u_{1,2}^\nu(t, x) - u_{2,1}^\nu(t, x)| \leq \varepsilon_\nu$$

on $[0, T] \times N$.

Now we know that $u_{1,2}^\nu$ converges to a variational solution of

$$\begin{cases} \frac{\partial}{\partial t_1} \hat{v}(t_1, t_2, x) + H_1(t_1, t_2, x, \frac{\partial}{\partial x} \hat{v}(t_1, t_2, x)) = 0 \\ \hat{v}(0, t_2, x) = \hat{w}(t_2, x) \end{cases}$$

where \hat{w} is the variational solution of the equation

$$\begin{cases} \frac{\partial}{\partial t_2} \hat{w}(t_2, x) + H_2(0, t_2, x, \frac{\partial}{\partial x} \hat{w}(t_2, x)) = 0 \\ \hat{w}(0, x) = f(x) \end{cases}$$

on the other hand, $u_{2,1}^\nu$ converges to a variational solution of

$$\begin{cases} \frac{\partial}{\partial t_2} \check{v}(t_1, t_2, x) + H_2(t_1, t_2, x, \frac{\partial}{\partial x} \check{v}(t_1, t_2, x)) = 0 \\ \check{v}(t_1, 0, x) = \check{w}(t_1, x) \end{cases}$$

where \check{w} is the variational solution of the equation

$$\begin{cases} \frac{\partial}{\partial t_1} \check{w}(t_1, x) + H_1(t_1, 0, x, \frac{\partial}{\partial x} \check{w}(t_1, x)) = 0 \\ \check{w}(0, x) = f(x) \end{cases}$$

Since we proved that $u_{1,2}^\nu(t, x) - u_{2,1}^\nu(t, x)$ goes to zero, the sequences have a common limit, v , and this will be a variational solution of both

$$\frac{\partial}{\partial t_2} v(t_1, t_2, x) + H_2(t_1, t_2, x, \frac{\partial}{\partial x} v(t_1, t_2, x)) = 0$$

and

$$\frac{\partial}{\partial t_1} v(t_1, t_2, x) + H_1(t_1, t_2, x, \frac{\partial}{\partial x} v(t_1, t_2, x)) = 0$$

and such that $v(0, 0, x) = f(x)$.

In other words, v is a variational solution of (MHJ). □

Remark 5.9. Note that the initial condition on the function u need not be given by “functions”. It is enough to have a “ C^0 - coisotropic submanifold” $C \subset T^*(\mathbb{R}^d \times N)$, that is the Hausdorff limit of submanifolds C_ν such that σ_{TC_ν} goes to zero for the C^0 topology, and $T^*(\mathbb{R}^d \times N) \rightarrow \mathbb{R}^d$ given by $(t_1, \dots, t_d, \tau_1, \dots, \tau_d, x, p) \rightarrow (t_1, \dots, t_d)$ when restricted to the leafs of the foliation is uniformly proper. We let the reader adapt such a generalization into our framework. We also mention that the notion of C^0 -Lagrangian submanifold has already been used by several authors, related to the fact that under various assumptions, we may guarantee that a smooth manifold that is the C^0 -limit of Lagrangian submanifolds has to be Lagrangian (see [Laudenbach-Sikorav-2],[Sikorav-3],[Viterbo 4]).

A Appendix: Proof of proposition 2.2

Lemma A.1. *Let $L_1, L_2, L_3 \in \mathcal{L}$. Then*

$$\gamma(L_1 - L_3) \leq \gamma(L_1 - L_2) + \gamma(L_2 - L_3)$$

Proof. Assume first that $L_2 = 0_N$. Then we must prove

$$\gamma(L_1 - L_3) \leq \gamma(L_1) + \gamma(L_3)$$

But this follows from corollary 2.8 and Proposition 3.3 on page 693 of [Viterbo 1] according to which

$$c(1, -S) = -c(\mu, S)$$

and

$$c(uv, S_1 + S_2) \geq c(u, S_1) + c(v, S_2)$$

Indeed, we have $\gamma(L) = \gamma(-L)$ and

$$c(1, S_1 - S_3) \geq c(1, S_1) + c(1, -S_3) = c(1, S_1) - c(\mu, S_3)$$

$$c(1, S_3 - S_1) \geq c(1, S_3) + c(1, -S_1) = c(1, S_3) - c(\mu, S_1)$$

using that $c(1, S_3 - S_1) = -c(\mu, S_1 - S_3)$ this can be rewritten as

$$-c(1, S_1 - S_3) \leq c(\mu, S_3) - c(1, S_1)$$

$$c(\mu, S_1 - S_3) \leq c(\mu, S_1) - c(1, S_3)$$

and adding the inequalities we get

$$\gamma(S_1 - S_3) \leq \gamma(S_1) + \gamma(S_3).$$

Then we claim that

$$(A.1) \quad \gamma(L - L') = \gamma(\varphi(L) - \varphi(L'))$$

for any φ in $\mathcal{H}\mathcal{D}$. Indeed, according to proposition 3.5 page 695 of [Viterbo 1], if $L' = \rho^{-1}(0_N)$, we have

$$\begin{aligned} \gamma(\varphi(L) - \varphi(L')) &= \gamma(\varphi(L) - \varphi\rho^{-1}(0_N)) = \\ \gamma((\varphi\rho^{-1})^{-1}(\varphi(L))) &= \gamma(\rho(L)) \end{aligned}$$

Since the last expression does not depend on the choice of φ , we get

$$\gamma(\varphi(L) - \varphi(L')) = \gamma(L - L')$$

Let now φ in $\mathcal{H}\mathcal{D}$ be such that $\varphi(L_2) = 0_N$, and let us apply the above equality to the L_j :

$$\begin{aligned} \gamma(L_1 - L_3) &= \gamma(\varphi(L_1) - \varphi(L_3)) \leq \gamma(\varphi(L_1)) + \gamma(\varphi(L_3)) = \\ \gamma(\varphi(L_1) - \varphi(L_2)) + \gamma(\varphi(L_2) - \varphi(L_3)) &= \gamma(L_1 - L_2) + \gamma(L_2 - L_3) \end{aligned}$$

□

We are now ready to prove proposition 2.3 stating the main properties of $\tilde{\gamma}$.

Proof of proposition 2.3. (a). Assume we have for all L in \mathcal{L} , $\gamma(\psi(L) - L) = 0$. According to [Viterbo 1], corollary 2.3, we have $\psi(L) = L$. Now since this holds for any L , we must have $\psi = \text{Id}$.

(b). Indeed the change of variable in \mathcal{L} given by $L = \psi(L')$ shows that

$$\tilde{\gamma}(\psi^{-1}) = \sup\{\gamma(\psi^{-1}(L) - L) \mid L \in \mathcal{L}\} = \sup\{\gamma(L' - \psi(L')) \mid L' \in \mathcal{L}\}$$

Since $\gamma(-S) = \gamma(S)$ we see that the last expression equals $\tilde{\gamma}(\psi)$

(c). Again by change of variable, we have

$$\begin{aligned} & \sup\{\gamma(\psi\varphi(L) - L) \mid L \in \mathcal{L}\} \leq \\ & \sup\{\gamma(\psi\varphi(L) - \varphi(L)) + \gamma(\varphi(L) - L) \mid L \in \mathcal{L}\} \leq \\ & \sup\{\gamma(\psi(L') - L') \mid L' \in \mathcal{L}\} + \sup\{\gamma(\varphi(L) - L) \mid L \in \mathcal{L}\} \end{aligned}$$

The second inequality follows from lemma A.1

(d). Indeed let us make the change of variable $L' = \varphi^{-1}(L)$, we get

$$\tilde{\gamma}(\varphi\psi\varphi^{-1}) = \sup\{\gamma((\varphi\psi\varphi^{-1})(L) - L) \mid L \in \mathcal{L}\} = \sup\{\gamma(\varphi\psi(L') - \varphi(L')) \mid L' \in \mathcal{L}\}$$

Now using the equality (A.1), we get

$$\gamma(\varphi\psi(L') - \varphi(L')) = \gamma(\psi(L') - L')$$

and thus

$$\sup\{\gamma(\varphi\psi(L') - \varphi(L')) \mid L' \in \mathcal{L}\} = \sup\{\gamma(\psi(L') - L') \mid L' \in \mathcal{L}\} = \tilde{\gamma}(\psi)$$

□

B Appendix : Symplectic invariants and the main theorem in the non-compact case

The aim of this appendix is to extend our construction and results to the case where N is a non-compact manifold. Of course, if all our Hamiltonians are compact supported, the statements and proofs are straightforward adaptations of the compact case. We refer to [Barles] and [Fathi-Maderna] for the study, from a different point of view, of Hamilton-Jacobi equations in the non-compact setting.

Let us first consider the problem of solving a single equation in our variational framework. For this we start from the isotropic manifold Λ_f , and consider $L =$

$\bigcup_{t \in \mathbb{R}} \Phi^t(\Lambda_{f,H})$. For this to be defined, we need that the Hamiltonian flow is complete, or at least that its restriction to $\Lambda_{0,H}$ is complete⁹

For a general Hamiltonian, we cannot hope to define variational solutions unless the associated flow is complete, but this condition will not suffice, as the following example shows.

Consider the equation

$$\begin{cases} \frac{\partial u}{\partial t} - x^2 - \frac{1}{4} \left| \frac{\partial u}{\partial x} \right|^2 = 0 \\ u(0, x) = 0 \end{cases}$$

The variational solution will be $u(t, x) = \tan(t)x^2$ which goes to infinity as t goes to $\frac{\pi}{2}$.

This happens because the Lagrangian submanifold associated to the equation is given by

$$L = \{(t, \tau, x, p) \mid \exists a \in \mathbb{R}, x = a \cos t, p = 2a \sin t, \tau = a^2\}$$

and this being vertical for $t = \frac{\pi}{2}$, cannot contain the graph of the differential of some function.

We see that the trouble comes from the fact that the flow at time $t = \frac{\pi}{2}$ brings over $\{x = 0\}$ points which are arbitrarily far. To avoid this phenomenon we introduce the notion of “finite propagation speed Hamiltonian”. We shall see that such a flow can be suitably approximated by compact supported flows, and such approximation can be used to define the variational solutions of the associated Hamilton-Jacobi equation.

We denote by $\pi : T^*N \rightarrow N$ the projection, by \mathcal{W} some exhaustion $(W_\mu)_{\mu \in \mathbb{N}}$ of T^*N by neighborhoods of the zero section, and by $(\Omega_\mu)_{\mu \in \mathbb{N}}$ some exhaustion¹⁰ of N .

We shall assume that for all μ , and all compact subsets K in N , $W_\mu \cap \pi^{-1}(K)$, is compact.

Our first task will be to determine when a sequence (φ_μ^t) of **compact supported** Hamiltonian isotopies **approximates** a given Hamiltonian isotopy, ψ^t , from the variational solution point of view. This will be based on

Proposition B.1. *Let σ^t be a compact supported Hamiltonian isotopy, $L = \sigma^1(0_N)$, ρ^t another compact supported Hamiltonian isotopy generated by $H(t, x, p)$, and set $\rho = \rho^1$. Assume we have $|H(t, x, p)| \leq \varepsilon$ on*

$$\mathcal{U}((\rho^t)_{t \in [0,1]}, L, \Omega) = \{(t, \rho^t(z)) \mid t \in [0, 1], z \in L, \rho(z) \in \pi^{-1}(\Omega)\}$$

⁹This means that $\Phi^s(z)$ is defined for all z in $\Lambda_{f,H}$.

¹⁰By exhaustion of a set X , we mean a non-decreasing sequence $(Y_\mu)_{\mu \in \mathbb{N}}$ of subsets of X such that $\bigcup_{\mu \in \mathbb{N}} Y_\mu = X$.

for some open set Ω in N .

Then for all x in Ω we have

$$|u_{\rho(L)}(x) - u_L(x)| \leq \varepsilon$$

In particular for $\varepsilon = 0$, this means that if H vanishes on

$$\bigcup_{t \in [0,1]} \{t\} \times \{\rho^t(L \cap \rho^{-1}(\pi^{-1}(\Omega)))\}$$

we have

$$u_{\rho(L)}(x) = u_L(x) \text{ for } x \text{ in } \Omega.$$

Proof. We first assume $|H| \leq \varepsilon$ everywhere. Then according to Proposition 2.6 we have $\gamma(\rho(L) - L) \leq \varepsilon$ and the inequality $|c(1_x, L') - c(1_x, L)| \leq \gamma(L' - L)$ implies

$$|u_{\rho(L)}(x) - u_L(x)| \leq \gamma(\rho(L) - L) \leq \varepsilon.$$

Now to deal with the general case, assume $(H_\tau)_{\tau \in [0,1]}$ is a smooth one-parameter family of compact supported Hamiltonians with $H_0 = H$, and for all τ in $[0, 1]$

$$H_\tau = H_0 \text{ on } \mathcal{U}(\rho^t, L, \Omega).$$

We claim that $c(1_x, \rho_\tau(L)) = c(1_x, L)$.

Now notice that $c(1_x, \rho_\tau(L))$ belongs to the set $C_\tau(x, L)$ of critical values of

$$(x, p, \xi) \longrightarrow \int_0^1 [p(s)\dot{x}(s) - H_\tau(s, x(s), p(s))] ds + S(x(0), \xi)$$

where S is a generating function quadratic at infinity for L .

The set $C_\tau(x, L)$ is equal to the set of values of

$$\int_0^1 [p_\tau(s)\dot{x}_\tau(s) - H_\tau(s, x_\tau(s), p_\tau(s))] ds + F(x_\tau(0), p(0))$$

where

$$(a) \quad (x_\tau(s), p_\tau(s)) = \rho_\tau^s(x_\tau(0), p_\tau(0))$$

$$(b) \quad (x_\tau(0), p_\tau(0)) \in L$$

$$(c) \quad x_\tau(1) = x$$

$$(d) \quad dF = \lambda \text{ on } L$$

Since by assumption, $z \in L$ and $\rho(z) \in \pi^{-1}(\Omega)$ imply $H_\tau = H_0$ on $\{(s, \rho^s(z)), s \in [0, 1]\}$, we conclude $\rho_\tau^s(z) = \rho^s(z)$.

Thus for x in Ω , the set $C_\tau(x, L)$ does not depend on τ . Since $C_\tau(x, L)$ is the set of critical values of the action, it has measure zero, and using the fact that $u_{\rho_\tau(L)}(x) = c(1_x, \rho_\tau(L))$ depends continuously on τ , we conclude that it must be independent of τ for x in Ω .

From the above argument follows that

$$u_{\rho_\tau(L)}(x) = u_{\rho(L)}(x) \text{ for all } x \text{ in } \Omega .$$

Now if $|H_0| \leq \varepsilon$ on $\mathcal{U}((\rho^t)_{t \in [0,1]}, L, \Omega)$ we may find a linear homotopy

$$H_\tau(s, z) = (1 - \tau\chi(s, z))H_0(s, z)$$

where $\chi(s, z) = 0$ for $(s, z) \in \mathcal{U}((\rho^t), L, \Omega)$ and $\chi = 1$ outside a neighborhood of $\mathcal{U}((\rho^t), L, \Omega)$.

By suitably choosing χ , we may assume the following inequality holds

$$|H_1| \leq \varepsilon + \delta$$

with $\delta > 0$ arbitrarily small. Since $H_\tau = H_0$ on $\mathcal{U}((\rho^t), L, \Omega)$, we have

$$u_{\rho_1(L)}(x) = u_{\rho(L)}(x)$$

for x in Ω . Since $|H_1| \leq \varepsilon + \delta$ everywhere, we may apply the first part of the proof and obtain

$$|u_{\rho_1(L)}(x) - u_L(x)| \leq \varepsilon + \delta$$

and thus

$$|u_{\rho(L)}(x) - u_L(x)| \leq \varepsilon + \delta .$$

Finally, δ being arbitrarily small, we conclude that

$$|u_{\rho(L)}(x) - u_L(x)| \leq \varepsilon$$

□

Our next task is to associate to the non compact supported Hamiltonian isotopy ψ^t a sequence $(\varphi_\mu^t)_{\mu \in \mathbb{N}}$ of compact supported ones, such that $c(1_x, \varphi_\mu^t(L))$ converges -or rather its restriction to any compact set stabilizes- for fixed x, t , as μ goes to infinity.

We denote by $H_\mu(t, z)$ and $H(t, z)$ the Hamiltonians generating φ_μ^t and ψ^t , and by $K_{\mu,\nu}(t, z)$ the Hamiltonian generating $\rho_{\mu,\nu}^t = \varphi_\nu^t \varphi_\mu^{-t}$.

The equality $\varphi_\mu^t = \varphi_\nu^t = \psi^t$ on \mathcal{U} is equivalent to $H_\mu(t, z) = H_\nu(t, z) = H(t, z)$ on $\psi^s(\mathcal{U})$ for all $s \in [0, T]$.

Thus if $\varphi_\mu^s = \varphi_\nu^s = \psi^s$ on

$$W_\mu \cap (\psi^t)^{-1}(\pi^{-1}(\Omega_\mu))$$

we have.

$$\varphi_\nu^s \varphi_\mu^{-s} = id \text{ on } \psi^s(W_\mu \cap (\psi^t)^{-1} \pi^{-1}(\Omega_\mu))$$

hence

$$K_{\mu,\nu}(s, z) = 0 \text{ on } \psi^s(W_\mu \cap (\psi^t)^{-1} \pi^{-1}(\Omega_\mu))$$

so that, according to proposition B.1,

$$u_{\varphi_\mu^t(L)}(x) = u_{\varphi_\nu^t(L)}(x) \text{ for } x \text{ in } \pi^{-1}(\Omega_\mu) \text{ } t \in [0, T]$$

This motivates the

Definition B.2. Let $(\psi^t)_{t \in [0, T]}$ be a Hamiltonian isotopy. A sequence of **compact supported** Hamiltonian isotopies, $(\varphi_\mu^t)_{\mu \in \mathbb{N}}$ is an **exhausting sequence** associated to \mathcal{W} and $(\Omega_\mu)_{\mu \in \mathbb{N}}$ if for all μ , we have

$$\varphi_\mu^t = \psi^t \text{ on } \bigcup_{s \in [0, T]} W_\mu \cap (\psi^s)^{-1}(\pi^{-1}(\Omega_\mu))$$

The above argument implies

Proposition B.3. Let (φ_μ^t) be an exhausting sequence for $(\psi^t)_{t \in [0, T]}$ associated to \mathcal{W} and $(\Omega_\mu)_{\mu \in \mathbb{N}}$. Then for any L , image of the zero section by a compact supported Hamiltonian isotopy and contained in some W_μ , and, for all $\alpha, \beta \geq \mu$ and $t \in [0, T]$

$$u_{\varphi_\alpha^t(L)}(x) = u_{\varphi_\beta^t(L)}(x) \quad \forall x \in \Omega_\mu$$

We may therefore set

Definition B.4. Given $(\psi^t)_{t \in [0, T]}$ and an exhausting sequence (φ_μ^t) associated to \mathcal{W} , $(\Omega_\mu)_{\mu \in \mathbb{N}}$ we define $u_{\psi^t(L)}(x)$ to be the common value of the $u_{\varphi_\mu^t(L)}(x)$ for μ large enough. Since $u_{\psi^t(L)}$ is a solution of the Hamilton-Jacobi equation, we call it the **variational solution associated to the exhausting sequence**.

We now need to solve two difficulties.

First, it is not clear, given $(\psi^t)_{t \in [0, T]}$, \mathcal{W} , $(\Omega_\mu)_{\mu \in \mathbb{N}}$ whether we may find an associated exhausting sequence. Second, one could wonder whether $u_{\psi^t(L)}(x)$ will depend on the choice of the exhausting sequence. But this will not be the case, since if $\check{\varphi}_\mu^t, \hat{\varphi}_\mu^t$ are exhausting sequences, we get a new exhausting sequence by intertwining them. Remember that we assume $W_\mu \cap \Omega_\mu$ is compact for all μ .

Definition B.5. (a). We shall say that $(\psi^t)_{t \in [0, T]}$ has **finite propagation speed** with respect to \mathcal{W} if and only if for each μ , there exists ν such that

$$\bigcup_{t \in [0, T]} \psi^t(W_\mu) \subset W_\nu$$

(b). \mathcal{W} is a **convex exhaustion** if the fibers of W_μ are convex, that is for all $x \in N$ $W_\mu \cap \pi^{-1}(x)$ is convex in T_x^*N .

Note that the composition of finite propagation speed Hamiltonians has finite propagation speed (with respect to the same family \mathcal{W}). However this is not necessarily true for its inverse $((\psi^t)^{-1})_{t \in [0, T]}$.

The set of Hamiltonians isotopies having, together with their inverse, finite propagation speed, is a group for the composition law

$$(\varphi^t) \circ (\psi^t) = (\varphi^t \psi^t)$$

Our last result for this section is

Proposition B.6. *Let \mathcal{W} be an exhaustion. If $(\psi^t)_{t \in [0, T]}$ has finite propagation speed with respect to \mathcal{W} , then, for any exhaustion $(\Omega_\mu)_{\mu \in \mathbb{N}}$ of N , there exists an exhausting sequence (φ_μ^t) associated to \mathcal{W} .*

The proof makes use of the obvious lemma:

Lemma B.7. *Let $K \subset U$ be a compact subset and $(\psi^t)_{t \in [0, 1]}$ a Hamiltonian isotopy such that*

$$V = \bigcup_{t \in [0, 1]} \psi^t(K) \subset U$$

Then there is a Hamiltonian isotopy, φ^t , supported near V such that

$$\varphi^t = \psi^t \text{ on } K \text{ for all } t \text{ in } [0, 1]$$

Proof. Indeed, if χ is a function equal to 1 on V , vanishing outside a neighborhood of U , and $H(t, z)$ generates ψ^t , then $\chi(z)H(t, z)$ generates φ^t . \square

Proof. We may now prove the proposition.

Since for all μ there exists ν such that

$$\psi^s(W_\mu) \subset W_\nu$$

for all s in $[0, T]$, we have

$$\psi^s(W_\mu) \cap \pi^{-1}(\Omega_\mu) \subset W_\nu \cap \pi^{-1}(\Omega_\mu) \subset W_\nu \cap \pi^{-1}(\Omega_\nu)$$

Thus

$$\bigcup_{s \in [0, T]} W_\mu \cap (\psi^s)^{-1}(\pi^{-1}(\Omega_\mu)) = \bigcup_{s \in [0, T]} (\psi^s)^{-1}(\psi^s(W_\mu) \cap \pi^{-1}(\Omega_\mu)) \subset \bigcup_{s \in [0, T]} (\psi^s)^{-1}(K_\nu)$$

We denote by $\tilde{K}_{\nu, T}$ this last set, and note that it is compact. Therefore

$$\bigcup_{s \in [0, T]} \psi^s(\tilde{K}_{\nu, T})$$

and we may find an isotopy φ_μ^t , compact supported, such that $\varphi_\mu^t = \psi^t$ on $\tilde{K}_{\nu, T}$. \square

Now this yields solutions of the evolution Hamilton-Jacobi equations in case the initial condition f has compact support. Of course formally the general case can be reduced to this one, since solving

$$\begin{cases} \frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) = 0 \\ u(0, x) = f(x) \end{cases}$$

is equivalent to solving

$$\begin{cases} \frac{\partial v}{\partial t}(t, x) + H(t, x, \frac{\partial v}{\partial x}(t, x)) + df(x) = 0 \\ v(0, x) = 0 \end{cases}$$

by the change of function $v(t, x) = u(t, x) - f(x)$ (and this change of function preserves variational solutions).

However we would like to have a criterion for H and an independent one for f in order to isolate the difficulties.

Proposition B.8. *Let \mathcal{W} be a convex exhaustion. Let $C_{\mathcal{W}}(N)$ be the union over all sets W_{μ} of the exhaustion, of the C^0 closure of the set of C^1 functions with graph in W_{μ} . Let $\mathcal{H}_{\mathcal{W}}^1(\mathbb{R} \times T^*N)$ be the set of $C^{1,1}$ Hamiltonians having finite propagation speed with respect to \mathcal{W} . Let f such that the graph of df is in W_{μ} , the function $u_{\psi^t(L)}$ does not depend on the choice of the exhausting sequence. Moreover it is a solution of the Hamilton-Jacobi equation*

$$(HJ) \quad \begin{cases} \frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) = 0 \\ u(0, x) = f(x) \end{cases}$$

Remark B.9. Let ψ^t be the flow of H and $\Lambda_0 \subset T^*N$ be a Lagrangian submanifold.

We first notice that the manifold we constructed, L in $T^*(\mathbb{R} \times N)$ giving the solution of (HJ) and the family $L_t = \psi^t(L)$ give essentially the same information. Indeed, if $u(t, x) = c(1_{(t,x)}, L)$ and $v(t, x) = c(1_x, L_t)$ we have $u(t, x) - v(t, x) = d(t)$ for some continuous function d . Given v , we can easily determine d or u , since $\frac{\partial}{\partial x}v(t, x) = \frac{\partial}{\partial x}u(t, x)$, hence

$$\frac{\partial}{\partial x}u(t, x) = -H(t, x, \frac{\partial}{\partial x}v(t, x))$$

hence

$$u(t, x) = f(x) + \int_0^t H(s, x, \frac{\partial}{\partial x}v(s, x))ds$$

Proof. Clearly if Λ_0 is the graph of df , we wish to replace it by a Lagrangian manifold Λ_{μ} , such that there exists an isotopy Λ_{μ}^s from Λ_0 to Λ_{μ} , such that

$$\varphi_{\mu}^t(\Lambda_{\mu}^s) = \psi^t(\Lambda_{\mu}^s) = \psi^t(\Lambda_0)$$

over $\pi^{-1}(\Omega_{\mu})$ for all $t \in [0, T]$.

But $\varphi_{\mu}^t = \psi^t$ on $W_{\mu} \cap (\psi^s)^{-1}\pi^{-1}(\Omega_{\mu})$ so if $\Lambda_{\mu}^s = \Lambda_0$ over $\pi^{-1}(A_{\mu})$, where A_{μ} will be determined later, we have

$$\varphi_{\mu}^t(\Lambda_{\mu}^s) = \psi^t(\Lambda_0)$$

over

$$\psi^t(W_{\mu} \cap (\psi^s)^{-1}\pi^{-1}(\Omega_{\mu}) \cap \pi^{-1}(A_{\mu}))$$

By compactness of

$$W_{\mu} \cap (\psi^s)^{-1}\pi^{-1}(\Omega_{\mu})$$

we may find A_{μ} such that $W_{\mu} \cap (\psi^s)^{-1}\pi^{-1}(\Omega_{\mu}) \subset \pi^{-1}(A_{\mu})$, and therefore according to proposition B.3

$$u_{\varphi_{\mu}^t(\Lambda_{\mu}^1)} = u_{\psi^t(\Lambda_{\mu}^1)}$$

on Ω_μ .

Now if Λ_0 is the graph of df , we may take for Λ_μ^s the graph of df_μ^s where

$$f_\mu^s(x) = (1 - s\chi_\mu(x))f(x)$$

and

$$\begin{cases} \chi_\mu(x) = 0 \text{ on } A_\mu \\ \chi_\mu(x) = 1 \text{ near infinity} \end{cases}$$

Then Λ_μ^s will coincide with Λ_0 over A_μ . We still need to prove that Λ_μ^s is in W_μ for all s in $[0, 1]$. Now this coincides with the graph of $(1 - s\chi_\mu(x))df(x)$ outside a compact set. Since the graph of df is in W_μ , this will be also in some W_ν for some ν , provided W_ν is fiber convex. □

Remark B.10. In practice most W_μ are fiber convex, and if this is not the case, we can always convexify them. However, it is not clear that a ψ^t having finite propagation speed with respect to \mathcal{W} will have also finite propagation speed with respect to the convexified exhaustion \mathcal{W}' , unless the convexification of W_μ , is contained in some W_ν .

Example B.11. (a). The Barles-Tourin condition defined for an autonomous Hamiltonian $H(x, p)$ on $T^*(\mathbb{R}^n)$ can be rewritten as

- $|D_p H(x, p)| \leq f(|p|)(1 + |x|)$

- $g_-(|p|) \leq |H(x, p)| \leq g_+(|p|)$

where $\lim_{\xi \rightarrow \infty} g_\pm(\xi) = +\infty$

This implies finite propagation speed for the exhaustion by the $U_r = \{(x, p) \mid |p| \leq r\}$, since the assumptions and conservation of energy imply that the image of $U_r = \{(x, p) \mid |p| \leq r\}$ by the flow remains in some $U_{r'}$. In such a region, the first inequality implies, by Gronwall's lemma, finite propagation speed.

(b). Consider the $U_{(a,b)} = \{(x, p) \mid |p| \leq ad(x, x_0) + b\}$, and assume

- $|D_x H(t, x, p)| \leq C(|p|)$

•

$$|D_p H(t, x, p)| \leq C(1 + d(x, x_0))$$

Then the first assumption, together with Gronwall's lemma implies that the flow sends $U_{a,b}$ to some $U_{a',b'}$.

The proposition implies that if \mathcal{W} is an exhaustion, and f a function on N such that the graph of df is contained in W_μ for some μ , and if $H(t, x, p)$ generates a flow having finite propagation speed with respect to the \mathcal{W} , then we may find a variational solution of

$$\begin{aligned} \frac{\partial u}{\partial t}(t, x) + H(t, x, \frac{\partial u}{\partial x}(t, x)) &= 0 \\ u(0, x) &= f(x) \end{aligned}$$

by considering solutions of

$$\begin{cases} \frac{\partial u_\mu}{\partial t}(t, x) + H_\mu(t, x, \frac{\partial}{\partial x} u_\mu(t, x)) = 0 \\ u_\mu(0, x) = f(x) \end{cases}$$

where H_μ is a compact supported Hamiltonian generating an exhausting sequence $(\varphi_\mu)_{\mu \in \mathbb{N}}$ for the flow ψ^t of H .

It follows from the above that $u(t, x) = \lim_{\mu \rightarrow \infty} u_\mu(t, x)$ defines a function independent of the choice of the exhausting sequence. In the regular case (i.e. for smooth data), this will of course be a solution of the equation almost everywhere.

We may of course combine theorem 5.6 with the above proposition to get the following:

Theorem B.12. *Let \mathcal{W} denote an exhausting sequence of neighborhoods of 0_N . Let $\mathfrak{C}_\mathcal{W}(N)$ be the union over all sets W_μ of the exhaustion, of the γ closures of the set of C^1 functions with graph in W_μ . Let $\mathcal{H}_\mathcal{W}^1(\mathbb{R} \times T^*N)$ be the set of $C^{1,1}$ Hamiltonians having finite propagation speed with respect to \mathcal{W} . Consider the closure $\mathfrak{H}_{\gamma, \mathcal{W}}(\mathbb{R} \times T^*N)$ of $\mathcal{H}_\mathcal{W}^1(\mathbb{R} \times T^*N)$ in $\mathfrak{H}_\gamma(\mathbb{R} \times T^*N)$. Finally we assume that*

$$(f, H_1, \dots, H_d) \in \mathfrak{C}(N) \times \mathfrak{H}_{\gamma, \mathcal{W}}(\mathbb{R} \times T^*N)^d$$

then there is a solution of (MHJ) with data (f, H_1, \dots, H_d) .

C Appendix : equations involving the unknown function

For equations of the type

$$\frac{\partial u}{\partial t_j} + H_j(t_1, \dots, t_d, x, \frac{\partial}{\partial x} u(x), u(x)) = 0$$

the same proofs hold, but we have to replace the cotangent space T^*M which is symplectic by the jet space $J^1(M)$, a contact manifold, on which we use the contact form $\alpha = dz - p dx$. We also replace Lagrangian submanifolds, by Legendrian ones, which are manifolds of dimension $\dim(M)$ on which α vanishes. It was noticed by Chekanov (around 1986, but published in [Chekanov] much later), that the theory of generating functions holds also in the Legendrian case, and that existence is invariant by Legendrian isotopy.

Since the notion of generating function applies “**even better**” to a Legendrian submanifold of $J^1(M)$, and according to [Théret], we also have uniqueness of generating functions, all the constructions of [Viterbo 1] apply. Similarly, since the Hamiltonian flow of $\tau + H(t, x, u, p)$ is also well-defined, the constructions from [Viterbo-Ottolenghi] also carry through.

The details of proofs are left to the reader. However, we will make explicit the commutativity condition for $H_1(t_1, t_2, x, p, u)$ and $H_2(t_1, t_2, x, p, u)$.

This is easy even in a general contact manifold. Indeed, define the Reeb vector field R , by the equalities

$$\begin{aligned} i_R \alpha &= 1 \\ i_R d\alpha &= 0 \end{aligned}$$

the associated Hamiltonian vector field must satisfy

$$\begin{aligned} i_{\tilde{X}_H} \alpha &= -H \\ i_{\tilde{X}_H} d\alpha &= dH + i_R(dH)\alpha \end{aligned}$$

(note that for $H = -1$ we recover R) which in our case, since $\alpha = du - p dx$, $R = \frac{\partial}{\partial u}$ translates to

$$\tilde{X}_H(x, p, u) = (pH_p - H) \frac{\partial}{\partial u} + H_p \frac{\partial}{\partial x} - (H_x + pH_u) \frac{\partial}{\partial p}$$

We then have $L_{\tilde{X}_H} \alpha = H_u \alpha$ and this shows that \tilde{X}_H preserves the contact distribution $\alpha = 0$. Since the commutator of such vector fields must also preserve the contact distribution, we may define

Definition C.1. Let $H_1(x, p, u)$ and $H_2(x, p, u)$ be two functions on $J^1(M)$. We set

$$\begin{aligned} & [H_1, H_2](x, p, u) = \\ & \{H_1, H_2\}_s(x, p, u) + \\ & \frac{\partial H_2}{\partial u}(x, p, u) \left(p \frac{\partial H_1}{\partial p}(x, p, u) - H_1(x, p, u) \right) - \frac{\partial H_1}{\partial u}(x, p, u) \left(p \frac{\partial H_2}{\partial p}(x, p, u) - H_2(x, p, u) \right) \end{aligned}$$

where

$$\{H_1, H_2\}_s = \frac{\partial H_1}{\partial p} \frac{\partial H_2}{\partial x} - \frac{\partial H_1}{\partial x} \frac{\partial H_2}{\partial p}$$

Proposition C.2.

$$\tilde{X}_{[H_1, H_2]} = [\tilde{X}_{H_1}, \tilde{X}_{H_2}]$$

Now in the multi-time dependent case, for $H_1(t_1, t_2, x, p, u)$, $H_2(t_1, t_2, x, p, u)$, we can consider

$$K_1(t_1, t_2, \tau_1, \tau_2, x, p, u) = \tau_1 + H_1(t_1, t_2, x, p, u), K_2(t_1, t_2, \tau_1, \tau_2, x, p, u) = \tau_2 + H_2(t_1, t_2, x, p, u)$$

as functions on $J^1(\mathbb{R}^2 \times M)$ and set

Definition C.3. Let H_1, H_2 as above. We define

$$[[H_1, H_2]] = [K_1, K_2]$$

The following is the natural extension of Theorem 1.2 :

Proposition C.4. Assume we have sequences H_ν, K_ν such that $H_\nu \xrightarrow{C^0} H, K_\nu \xrightarrow{C^0} K, [H_\nu, K_\nu] \xrightarrow{C^0} 0$. Then $[H, K] = 0$.

D Appendix : Coisotropic submanifolds of symplectic manifolds

Let (P, σ) be a symplectic manifold, and C a coisotropic submanifold, that is

$$(T_z C)^\perp \subset T_z C \quad \forall z \in C$$

where

$$(T_z C)^\perp := \{v \in T_z P : \sigma(v, u) = 0 \quad \forall u \in T_z C, \}$$

here σ is a closed and non degenerated 2-form, e.g. if $P = T^*M$, then $\sigma = d\theta$, $\theta = \sum_i p_i dx^i$ is the Liouville 1-form. Let $H : T^*M \rightarrow \mathbb{R}$, then the Hamiltonian vector field X_H related to H is so defined:

$$\sigma(X_H, \cdot) = -dH,$$

and the Poisson brackets of two Hamiltonian functions H e K are

$$\{K, H\}(z) = dK(z) \cdot X_H(z) \quad (z = (x, p)).$$

Let suppose that C is defined by zero's of functions¹¹, $H^a : P \rightarrow \mathbb{R}$, $a = 1, \dots, \text{codim } C$:

$$C : H^a = 0 \quad \text{rk } dH^a|_{H^a=0} = \max = \text{codim } C$$

Thus: $u \in T_z C \iff dH^a u = 0$. Standard homomorphism theorem between linear spaces gives existence (and uniqueness, by the max rank) of the Lagrange multipliers $\lambda_a = \lambda_a(z, v)$:

$$\forall v \in (T_z C)^\perp : [dH^a u = 0 \Rightarrow \sigma(v, u) = 0] \Rightarrow \omega(v, \cdot) = \sum_a \lambda_a dH^a$$

The linear independence of the forms dH^a , $a = 1, \dots, \text{codim } C$, implies that the following Hamiltonian vector fields $(X_a)_{a=1, \dots, \text{codim } C}$,

$$\sigma(X_a, \cdot) = -dH^a,$$

span $(T_z C)^\perp$. Since $(T_z C)^\perp \subset T_z C$, the vectors $X_a(z)$ are in $T_z C$, and their integral paths, the *characteristics* of X_a , are in C , hence

$$\sigma(X_a, X_b) = 0.$$

From the definition of Poisson brackets $\{\cdot, \cdot\}$,

$$\sigma(X_a, X_b) = \{H_a, H_b\},$$

and from the well known Lie algebra morphism $([\cdot, \cdot]: \text{Lie brackets})$

$$[X_a, X_b] = X_{\{H_a, H_b\}},$$

one obtains that the fields $(X_a)_{a=1, \dots, \text{codim } C}$ are in *involution*:

$$[X_a, X_b] = 0.$$

In other words, by Frobenius' theorem, $(TC)^\perp$ is a *integrable distribution*, the *characteristic* distribution. Notice that the involution condition of the Hamiltonians

$$\{H_a, H_b\} = 0$$

is intrinsic, constitutive, of the coisotropic submanifold C .

¹¹it is equivalent to ask that the normal bundle of C , with respect to some Riemannian metric, is trivial ([Guillemin-Pollack] p. 77.)

E Appendix: Geometric Theory of Hamilton-Jacobi equations on coisotropic submanifolds

Let us consider d functionally independent and in involution Hamiltonians \tilde{H}_a , $a = 1, \dots, d$, of the following structure:

$$\begin{aligned} \tilde{H}_a : T^*(N \times \mathbb{R}^d) &\longrightarrow \mathbb{R} \\ (x_1, \dots, x_n, t_1, \dots, t_d, p_1, \dots, p_n, \tau_1, \dots, \tau_d) &\longmapsto \tau_a + H_a(t_1, \dots, t_d, x, p) \end{aligned}$$

The submanifold

$$C = \bigcap_{a=1}^d \tilde{H}_a^{-1}(0)$$

is coisotropic (because $\{\tilde{H}_a, \tilde{H}_b\} = 0$), $\dim C = 2n + d$, $\text{co-dim } C = d$.

We call *geometric solution* of the Hamilton-Jacobi multi-equation related to C every *Lagrangian submanifold*¹² $\Lambda \subset T^*(N \times \mathbb{R}^d)$ belonging to C :

$$\Lambda \subset C.$$

The Lagrangian submanifolds which are graphs of a section of the cotangent bundle $T^*(N \times \mathbb{R}^d)$ are image of the differential of functions S , defined on $N \times \mathbb{R}^d$, $S : N \times \mathbb{R}^d \rightarrow \mathbb{R}$,

$$\Lambda = \text{im } dS = \left\{ (x, t, p, \tau) : p = \frac{\partial S}{\partial x}(x, t), \tau = \frac{\partial S}{\partial t}(x, t) \right\}$$

Thus, every globally transverse Lagrangian submanifold Λ , solving the H-J multi-equation related to C , represents, by its generating function S , also a *classic* solution of it:

$$(A1) \quad \frac{\partial S}{\partial t_a}(x, t) + H_a(t_1, \dots, t_d, x, \frac{\partial S}{\partial x}(x, t)) = 0, \quad a = 1, \dots, d.$$

• The characteristics method. The following proposition suggests us how to build geometric solutions of the H-J multi-equation:

Proposition *Let $\Lambda \subset C$, Λ be Lagrangian and $C = \bigcap_{a=1}^d \tilde{H}_a^{-1}(0)$ coisotropic. Then the Hamiltonian vector fields $X_{\tilde{H}_a}$, $a = 1, \dots, d$, are tangent to Λ .*

Proof. If $v \in T_x \Lambda \Rightarrow v \in T_x C \Rightarrow v \in \bigcap_{a=1}^d \ker d\tilde{H}_a(x)$, that is $d\tilde{H}_a(x)v = 0$, $\forall a = 1, \dots, d$. It holds that $\sigma(X_{\tilde{H}_a}, v) = -d\tilde{H}_a v = 0$, that is, the fields $X_{\tilde{H}_a}$ are skew-orthogonal to Λ , $X_{\tilde{H}_a}(x) \in (T_x \Lambda)^\perp = T_x \Lambda$, where the last equality holds since Λ is Lagrangian. \square

¹²that is, *i*) $\sigma|_\Lambda = 0$ and *ii*) $\dim \Lambda = \dim N \times \mathbb{R}^d = n + d$.

• **Multi-valued Variational Solutions.** The Liouville 1-form of $T^*(N \times \mathbb{R}^d)$ is

$$\theta = \sum_{j=1}^n p_j dx^j + \sum_{a=1}^d \tau_a dt^a$$

The restriction (pull-back) of θ on the Lagrangian submanifolds gives closed 1-forms, indeed, by denoting j the inclusion map $j : \Lambda \hookrightarrow T^*(N \times \mathbb{R}^d)$,

$$0 = \sigma|_{\Lambda} = j^* \sigma = j^* d\theta = dj^* \theta = d(\theta|_{\Lambda}).$$

Let $f : N \rightarrow \mathbb{R}$ be the initial data for the multi-equation (A1), then a primitive function of $\theta|_{\Lambda}$ (on Λ , parametrized by χ, t^1, \dots, t^d) is given by

$$\bar{S}(\chi, t^1, \dots, t^d) = f(\chi) + \int_{\tilde{\Phi}^{t^1} \dots \tilde{\Phi}^{t^a} \dots \tilde{\Phi}^{t^d}(z) \Big|_{t^a \in [0, t^a]}} \sum_{j=1}^n p_j dx^j - \sum_{a=1}^d H_a dt^a,$$

$$z = (\chi, 0; df(\chi), -H_a(0, \chi, df(\chi))) \in T^*(N \times \mathbb{R}^d)$$

For small t^1, \dots, t^d , the x -components of $\tilde{\Phi}^{t^1} \dots \tilde{\Phi}^{t^a} \dots \tilde{\Phi}^{t^d}(z)$, i.e.

$$x(\chi, t) = \text{pr}_N \pi_{N \times \mathbb{R}^d} \tilde{\Phi}^{t^1} \dots \tilde{\Phi}^{t^a} \dots \tilde{\Phi}^{t^d}(z),$$

admits inverse: $\chi = \chi(x, t)$. In such a (scarcely meaningful) case, the function

$$S(x, t) := \bar{S}(\chi(x, t), t)$$

represents the classic solution to the Cauchy problem.

F Appendix: A short account on the developments of the theory of the multi-equations of Hamilton-Jacobi

After a pioneering paper by Tulczyjew [Tulczyjew], the study Hamilton-Jacobi problems in general co-isotropic submanifolds of co-dimension greater than one inside T^*Q , that is, extending the standard case $\frac{\partial S}{\partial t} + H(t, x, \frac{\partial S}{\partial x}) = 0$, was further developed by Benenti and Tulczyjew in [Benenti-Tulczyjew] and then in [Benenti], where, among other things, in this last paper first it was considered generating functions with finite auxiliary parameters for Lagrangian submanifolds of $T^*Q \times T^*Q$ representing symplectic relations (canonical transformations). Examples of this new framework are the integrable Dirac systems reconsidered by [Lichnerowicz] and

[Menzio-Tulczyjew]. More recently, new examples of H-J multi-equations arose from economics [Rochet], one can see the bibliography in [Barles-Tourin]. The use of symplectic topology and the notion of variational solution for the H-J equation have been introduced by Chaperon and Sikorav, and then developed in [Viterbo 2], where, in particular, the uniqueness of such a solution is deduced by the essential global uniqueness of the generating functions proved in [Viterbo 1]. We recall that these variational solutions coincide with the viscosity solutions in the case of p -convex Hamiltonians [Zhukovskaya], but it is not true in general case, lacking a (expected) Markov condition (cf. [Viterbo 2], [Viterbo-Ottolenghi]).

References

- [Barles] G. Barles, *Solutions de viscosité des équations de Hamilton-Jacobi*. Mathématiques et applications, vol. 17, Springer-Verlag, 1994.
- [Barles-Tourin] G. Barles, A. Tourin, Commutation properties of semigroups for first order Hamilton-Jacobi equations and application to multi-time equations, *Indiana Univ. Math. Journal*, vol 50, (2001), pp. 1524-1543.
- [Benenti] S. Benenti, *Symplectic relations in analytical mechanics*, Proceedings of the IUTAM-ISIMM symposium on modern developments in analytical mechanics, Vol. I (Torino, 1982). Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. 117 (1983), suppl. 1, 39–91.
- [Benenti-Tulczyjew] S. Benenti, W.M. Tulczyjew, *The geometric meaning and globalization of the Hamilton-Jacobi method*, Differential geometric methods in mathematical physics (Proc. Conf., Aix-en-Provence/Salamanca, 1979), pp. 9–21, Lecture Notes in Math., 836, Springer, Berlin, 1980.
- [Bernardi-Cardin] O. Bernardi, F. Cardin, Minmax and viscosity solutions of Hamilton-Jacobi equations in the convex case *Communications on Pure and Applied analysis* Volume: 5, Number: 4 (2006), pp. 793-812.

- [Borel] A. Borel, *Cohomologie des espaces localement compacts (d'après Leray)*. Springer Lecture Notes, vol. 2, Springer-Verlag, Berlin-Heidelberg-New-York.
- [Brunella] M. Brunella, On a theorem of Sikorav *Enseignement Mathématique* Volume:37,(1991), pp. 83–87.
- [Chaperon] M. Chaperon, Lois de conservation et géométrie symplectique. *C. R. Acad. Sci. Paris. Série I, Math.*, 312 (1991), pp. 345-348.
- [Chekanov] Yu. V. Chekanov, Critical points of quasifunctions, and generating families of Legendrian manifolds. *Funktsional. Anal. i Prilozhen.* vol. 30 (1996), pages 56–69. Translation in *Funct. Anal. Appl.* vol. 30, (1996), no. 2, pages 118–128.
- [Crandall-Lions] M.G. Crandall and P.-L. Lions. Viscosity solutions of Hamilton–Jacobi equations. *Trans. Amer. Math. Soc.*, 277:1–43, 1983.
- [Dacorogna-Marcellini] B. Dacorogna, P. Marcellini, *Implicit Partial Differential equations*. Birkhauser, Boston, 1999.
- [Ekeland-Hofer] I. Ekeland , H. Hofer, Symplectic topology and Hamiltonian dynamics. *Math. Z.* 200 (1990) 355-378.
- [Eliashberg] Y. Eliashberg, Rigidity for symplectic and contact structures. Preprint, 1981.
- [Entov-Polterovich-Zapolsky] M. Entov, L. Polterovich and F. Zapolsky, *Quasi-morphisms and the Poisson bracket*. math.SG/0605406
- [Fathi-Maderna] A. Fathi, E. Maderna, Weak KAM theorem on non compact manifolds. *Nolinear Differential Equations and Applications*, vol. 11, n. 3, p. 271-300, 2004.
- [Gromov] Misha Gromov. Pseudoholomorphic curves in symplectic manifolds. *Invent. Math.*, 82(2):307–347, 1985.
- [Guillemin-Pollack] V. Guillemin, A. Pollack, *Differential topology*, Prentice-Hall Inc., Englewood Cliffs, N.J., 1974.

- [Hofer] H. Hofer, On the topological properties of symplectic maps. *Proceedings of the Royal Society of Edinburgh*, vol.115 (1990), pp. 25-38.
- [Humilière] V. Humilière. *On some completions of the space of Hamiltonian maps*. math.SG/0511418 (submitted)
- [Humilière 2] V. Humilière. *Quasi representations of nilpotent Lie algebras are C^0 closed*. in preparation, 2006.
- [Laudenbach-Sikorav] F. Laudenbach and J.-C. Sikorav, Persistence d'intersection avec la section nulle au cours d'une isotopie hamiltonienne dans un fibré cotangent, *Inventiones Math.* vol. 82 (1985) pp. 349-357.
- [Laudenbach-Sikorav-2] François Laudenbach and Jean-Claude Sikorav. Hamiltonian disjunction and limits of lagrangian submanifolds. *Intern. Math. Res. Notes*, 4:161–168, 1994.
- [Lichnerowicz] A. Lichnerowicz, Variété symplectique et dynamique associée à une sous-variété. (French) *C. R. Acad. Sci., Paris, Sér. A* 280, 523-527, (1975).
- [Lions-Rochet] P.-L. Lions and J.-C. Rochet, *Hopf formulas and multi-time Hamilton-Jacobi equations*, *Proc. Amer. Math. Soc.* 96 (1986), pp. 79-84.
- [Motta-Rampazzo] Monica Motta and Franco Rampazzo, Nonsmooth multi-time hamilton-jacobi systems. *Indiana Univ. Math. J.*, 55:1573–1614, 2006.
- [Menzio-Tulczyjew] M.R. Menzio and W.M.Tulczyjew, Infinitesimal symplectic relations and generalized Hamiltonian dynamics. *Ann. Inst. Henri Poincaré, Nouv. Sér.*, Sect. A 28, 349-367 (1978).
- [Rampazzo-Sussmann] F. Rampazzo, H. Sussmann, *Set-valued differentials and a nonsmooth version of Chow's theorem*. Published in the Proceedings of the 40th IEEE Conference on Decision and Control; Orlando, Florida, December 4 to 7, 2001 (IEEE Publications, New York, 2001), Volume 3, pp. 2613-2618.

- [Rochet] J.-C. Rochet, The taxation principle and multi-time Hamilton-Jacobi Equation *J. of Math. Econ.* 14 (1985), 113-128.
- [Sikorav 1] J.-C. Sikorav, Personal communication. (1990).
- [Sikorav-2] J.-C. Sikorav, Problèmes d'intersections et de points fixes en géométrie hamiltonienne. *Comment. Math. Helv.* 62, No.1, pp. 62-73 (1987).
- [Sikorav-3] Jean-Claude Sikorav. Quelques propriétés des plongements lagrangiens. *Mém. S.M.F. n°46, suppl. au Bull; S.M.F. 119 (1991).*, pages 151–167, 1991.
- [Théret] D. Théret, Thèse de Doctorat *Université de Paris 7, Denis Diderot, 1996.*
- [Theret 2] D. Théret, *A complete proof of Viterbo's uniqueness theorem on generating functions.* *Topology Appl.* 96, no. 3, 249-266 (1999).
- [Tulczyjew] W. M. Tulczyjew, *Relations symplectiques et les équations d'Hamilton-Jacobi relativistes,* C. R. Acad. Sc. Paris 281 A, 545-547, 1975.
- [Viterbo 1] C. Viterbo, Symplectic topology as the geometry of generating functions, *Mathematische Annalen*, 292, (1992), pp. 685–710.
- [Viterbo 2] C. Viterbo, *Solutions d'équations de Hamilton-Jacobi*, Séminaire X-EDP 1995, Palaiseau.
- [Viterbo 3] C. Viterbo, Metric and isoperimetric problems in symplectic topology, *J. Amer. Math. Soc.* 13 (2000), no. 2, pp. 411–431.
- [Viterbo 4] C. Viterbo, On the uniqueness of generating Hamiltonian for continuous limits of Hamiltonians flows, *International Mathematics Research Notices 2006* Article ID 34028, 9 pages, 2006. doi:10.1155/IMRN/2006/34028.
- [Viterbo-Ottolenghi] C. Viterbo, A. Ottolenghi, *Solutions d'équations d'Hamilton-Jacobi et géométrie symplectique,*

Preprint, 1995

<http://math.polytechnique.fr/cmat/viterbo/Semi.HJ.dvi>.

[Weinstein]

A. Weinstein, *Lectures on symplectic geometry*,
CBMS Lecture Series, AMS, 1974.

[Zhukovskaya]

T. Zhukovskaya, *Singularités de minimax et solutions faibles d'Equations aux dérivées partielles*.
Thèse de Doctorat, Université de Paris 7, 1993.

[Zhukovskaya 2]

T. Zhukovskaya, *Metamorphoses of the Chaperon-Sikorav weak solutions of Hamilton-Jacobi equations*
Thèse de Doctorat, Université de Paris 7, 1993.