

QUANTUM CHARACTERS FOR QUANTUM AFFINE ALGEBRAS

DAVID HERNANDEZ

1. INTRODUCTION

In the case of semi-simple Lie algebras, the structure of the Grothendieck ring of finite dimensional representations of the quantum algebra is well understood: it is analogous to the classic case $q=1$ and we have a ring homomorphism of characters.

For the general case of Kac-Moody algebras the picture is less clear. In the affine case, E. Frenkel and N. Reshetikhin introduced an injective ring homomorphism of q -characters. It gives informations about the decomposition in Jordan subspaces for a class of commutative elements. The homomorphism of q -characters has a nice symmetry property analogous to the classic action of the Weyl group: the image is the intersection of the kernels of screening operators.

In the *ADE* case, H. Nakajima, motivated by the geometry of quiver varieties, introduced t -analogs of q -characters. The definition is combinatorial but the proof of the existence uses the geometric theory of quiver varieties which holds only in the simply laced case. In the preprint math.QA/0212257 we propose an algebraic general new approach to q, t -characters motivated by deformed screening operators. The t -deformations are naturally deduced from the algebra structure of $\mathcal{U}_q(\hat{\mathfrak{h}})$: the parameter t is analog to the central charge $c \in \mathcal{U}_q(\hat{\mathfrak{h}})$. This variant of Nakajima's theory allows us to treat the non-simply laced case: in particular the morphism of q, t -characters $\chi_{q,t}$ leads to the construction of a quantization of the Grothendieck ring and to general Nakajima's analogs of Kazhdan-Lusztig polynomials.

2. HOMOMORPHISM OF CHARACTERS IN FINITE CASE

Notations:

\mathfrak{g} : simple Lie algebra of rank n , $I = \{1, \dots, n\}$

$\mathfrak{h} \subset \mathfrak{g}$: Cartan subalgebra

$\Lambda \subset \mathfrak{h}^*$: lattice of weights of \mathfrak{g}

$\omega_i \in \Lambda$: fundamental weights

$\text{Rep}(\mathcal{U}(\mathfrak{g}))$: Grothendieck ring of finite dimensional representations (with \oplus and \otimes)

We have an injective homomorphism of rings:

$$\chi : \text{Rep}(\mathcal{U}(\mathfrak{g})) \rightarrow \mathbb{Z}[y_i^\pm]_{i \in I}$$

$$\chi(V) = \sum_{\lambda = \sum_{i \in I} m_i \omega_i \in \Lambda} \dim(V_\lambda) \prod_{i \in I} y_i^{m_i}$$

where V_λ is the weight space of V :

$$V_\lambda = \{x \in V / \forall h \in \mathfrak{h}, h.x = \lambda(h)x\}$$

We have a symmetry property related to the Weyl groups W :

$$\text{Im}(\chi) = \mathbb{Z}[y_i^\pm]_{i \in I}^W \simeq \mathbb{Z}[T_1, \dots, T_n]$$

The quantum case is analogueous:

$$\text{Rep}(\mathcal{U}_q(\mathfrak{g})) \simeq \text{Rep}(\mathfrak{g}) \simeq \mathbb{Z}[\Lambda]^W \simeq \mathbb{Z}[T_1, \dots, T_n]$$

where $\text{Rep}(\mathcal{U}_q(\mathfrak{g}))$: finite-dimensional representations of type 1.

In the quantum affine case $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$, the picture is less clear.

3. QUANTUM AFFINE ALGEBRA

\mathfrak{g} : simple Lie algebra of rank n , $I = \{1, \dots, n\}$

$(C_{ij})_{1 \leq i, j \leq n} = (\frac{2(\alpha_i, \alpha_j)}{(\alpha_i, \alpha_i)})$: Cartan matrix of \mathfrak{g}

$\mathfrak{g} \subset \hat{\mathfrak{g}}$: affine Lie algebra, $(C_{i,j})_{0 \leq i, j \leq n}$ generalized Cartan matrix of $\hat{\mathfrak{g}}$

$r_i = \frac{(\alpha_i, \alpha_i)}{2} \in \mathbb{Z}$, $B_{i,j} = r_i C_{i,j}$ symmetric matrix

$q \in \mathbb{C}^*$ not a root of unity, $q_i = q^{r_i}$

Definition 1. The quantum affine algebra $\mathcal{U}_q(\hat{\mathfrak{g}})$ is defined by generators $x_{i,m}^\pm$ ($1 \leq i \leq n$, $m \in \mathbb{Z}$), k_i^\pm ($1 \leq i \leq n$), $h_{i,m}$ ($1 \leq i \leq n$, $m \in \mathbb{Z}^*$), central elements $c^{\pm \frac{1}{2}}$ and relations:

$$\begin{aligned} k_i k_j &= k_j k_i \\ k_i h_{j,m} &= h_{j,m} k_i \\ k_i x_{j,m}^\pm k_i^{-1} &= q^{\pm B_{ij}} x_{j,m}^\pm \\ [h_{i,m}, x_{j,m'}^\pm] &= \pm \frac{1}{m} [m B_{ij}]_q c^{\mp \frac{|m|}{2}} x_{j,m+m'}^\pm \\ x_{i,m+1}^\pm x_{j,m'}^\pm - q^{\pm B_{ij}} x_{j,m'}^\pm x_{i,m+1}^\pm &= q^{\pm B_{ij}} x_{i,m}^\pm x_{j,m'+1}^\pm - x_{j,m'+1}^\pm x_{i,m}^\pm \\ [h_{i,m}, h_{j,m'}] &= \delta_{m,-m'} \frac{1}{m} [m B_{ij}]_q \frac{c^m - c^{-m}}{q - q^{-1}} \\ [x_{i,m}^+, x_{j,m'}^-] &= \delta_{ij} \frac{c^{\frac{m-m'}{2}} \phi_{i,m+m'}^+ - c^{-\frac{m-m'}{2}} \phi_{i,m+m'}^-}{q_i - q_i^{-1}} \\ \sum_{\pi \in \Sigma_s} \sum_{k=0 \dots s} (-1)^k \begin{bmatrix} s \\ k \end{bmatrix}_{q_i} x_{i,m_{\pi(1)}}^\pm \dots x_{i,m_{\pi(k)}}^\pm x_{j,m'}^\pm x_{i,m_{\pi(k+1)}}^\pm \dots x_{i,m_{\pi(s)}}^\pm &= 0 \end{aligned}$$

where the last relation holds for all $i \neq j$, $s = 1 - C_{ij}$, all sequences of integers m_1, \dots, m_s . Σ_s is the symmetric group on s letters. For $i \in I$ and $m \in \mathbb{Z}$, $\phi_{i,m}^\pm \in \mathcal{U}_q(\hat{\mathfrak{g}})$ is determined by the formal power series in $\mathcal{U}_q(\hat{\mathfrak{g}})[[u]]$ (resp. in $\mathcal{U}_q(\hat{\mathfrak{g}})[[u^{-1}]]$):

$$\sum_{m=0 \dots \infty} \phi_{i,\pm m}^\pm u^{\pm m} = k_i^\pm \exp(\pm(q - q^{-1}) \sum_{m'=1 \dots \infty} h_{i,\pm m'} u^{\pm m'})$$

and $\phi_{i,m}^+ = 0$ for $m < 0$, $\phi_{i,m}^- = 0$ for $m > 0$.

One has an embedding $\mathcal{U}_q(\mathfrak{g}) \subset \mathcal{U}_q(\hat{\mathfrak{g}})$ and a Hopf algebra structure on $\mathcal{U}_q(\hat{\mathfrak{g}})$.

4. FINITE DIMENSIONAL REPRESENTATIONS OF $\mathcal{U}_q(\hat{\mathfrak{g}})$

A finite dimensional representation V of $\mathcal{U}_q(\hat{\mathfrak{g}})$ is called of type 1 if c acts as Id and V is of type 1 as a representation of $\mathcal{U}_q(\mathfrak{g})$. We note $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ the Grothendieck ring of finite dimensional representations of type 1.

The operators $\{\phi_{i,\pm m}^\pm, i \in I, m \in \mathbb{Z}\}$ commute on V . So we have a pseudo weight space decomposition:

$$V = \bigoplus_{\gamma \in \mathbb{C}^I \times \mathbb{Z} \times \mathbb{C}^I \times \mathbb{Z}} V_\gamma$$

where for $\gamma = (\gamma^+, \gamma^-)$, V_γ is a simultaneous generalized eigenspace:

$$V_\gamma = \{x \in V / \exists p \in \mathbb{N}, \forall i \in \{1, \dots, n\}, \forall m \in \mathbb{Z}, (\phi_{i,m}^\pm - \gamma_{i,m}^\pm)^p \cdot x = 0\}$$

The $\gamma_{i,m}^\pm$ are called pseudo eigen values of V .

Theorem 1. (Chari, Pressley 94) *Every simple representation $V \in \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ is a highest weight representation V , that is to say there is $v_0 \in V$ (highest weight vector) $\gamma_{i,m}^\pm \in \mathbb{C}$ (highest weight) such that:*

$$V = \mathcal{U}_q(\hat{\mathfrak{g}}).v_0, \quad c^{\frac{1}{2}}.v_0 = v_0 \\ \forall i \in I, m \in \mathbb{Z}, x_{i,m}^+.v_0 = 0, \quad \phi_{i,m}^\pm.v_0 = \gamma_{i,m}^\pm v_0$$

Moreover we have an I -uplet $(P_i(u))_{i \in I}$ of (Drinfeld-)polynomials such that $P_i(0) = 1$ and:

$$\gamma_i^\pm(u) = \sum_{m \in \mathbb{N}} \gamma_{i,\pm m}^\pm u^\pm = q_i^{\deg(P_i)} \frac{P_i(uq_i^{-1})}{P_i(uq_i)} \in \mathbb{C}[[u^\pm]]$$

and $(P_i)_{i \in I}$ parametrizes simple modules in $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$.

Theorem 2. (Frenkel, Reshetikhin 98) *The eigenvalues $\gamma_i(u)^\pm \in \mathbb{C}[[u]]$ of a representation $V \in \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ have the form:*

$$\gamma_i^\pm(u) = q_i^{\deg(Q_i) - \deg(R_i)} \frac{Q_i(uq_i^{-1})R_i(uq_i)}{Q_i(uq_i)R_i(uq_i^{-1})}$$

where $Q_i(u), R_i(u) \in \mathbb{C}[u]$ and $Q_i(0) = R_i(0) = 1$.

Note that the polynomials Q_i, R_i are uniquely defined by γ . We note $Q_{\gamma,i}, R_{\gamma,i}$ the polynomials associated to γ .

Example: We suppose $\mathfrak{g} = \mathfrak{sl}_2$ and so $I = \{1\}$. We do explicit computations with the help of Jimbo's evaluation homomorphism $ev_a : \mathcal{U}_q(\hat{\mathfrak{sl}}_2) \rightarrow \mathcal{U}_q(\mathfrak{sl}_2)$.

For Drinfeld-polynomial $P(u) = 1 - ua$ we have the $\mathcal{U}_q(\hat{\mathfrak{sl}}_2)$ -module $M_a = \mathbb{C}v_0 \oplus \mathbb{C}v_1$:

$$v_0 \text{ is the highest weight vector: } \phi^\pm(u).v_0 = q^{\frac{1-uaq^{-1}}{1-uaq}} v_0 = q^{\frac{P(uq^{-1})}{P(uq)}} v_0.$$

v_1 is a simultaneous eigenvector:

$$\phi^\pm(u).v_1 = q^{-1} \frac{1-uaq^3}{1-uaq} v_1 = q^{-1} \frac{Q(uq^{-1})R(uq)}{Q(uq)R(uq^{-1})} v_1$$

where

$$Q(u) = (1 - ua), \quad R(u) = (1 - uaq^2)(1 - ua)$$

5. FRENKEL-RESHETIKHIN'S Q-CHARACTERS

Let \mathcal{Y} be the commutative ring $\mathcal{Y} = \mathbb{Z}[Y_{i,a}^\pm]_{i \in I, a \in \mathbb{C}^*}$.

Definition 2. For $V \in \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ a representation, the q -character $\chi_q(V)$ of V is:

$$\chi_q(V) = \sum_{\gamma \in \mathbb{C}^I \times \mathbb{Z} \times \mathbb{C}^I \times \mathbb{Z}} \dim(V_\gamma) \prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{\lambda_{\gamma,i,a} - \mu_{\gamma,i,a}} \in \mathcal{Y}$$

where

$$Q_{\gamma i}(z) = \prod_{a \in \mathbb{C}^*} (1 - za)^{\lambda_{\gamma,i,a}}, \quad R_{\gamma i}(z) = \prod_{a \in \mathbb{C}^*} (1 - za)^{\mu_{\gamma,i,a}}$$

Example: $\chi_q(M_a) = Y_a + Y_{aq^2}^{-1}$

Theorem 3. (Frenkel, Reshetikhin 98) *The map*

$$\chi_q : \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) \rightarrow \mathcal{Y}$$

is an injective ring homomorphism and the following diagram is commutative:

$$\begin{array}{ccc} \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) & \xrightarrow{\chi_q} & \mathbb{Z}[Y_{i,a}^\pm]_{i \in I, a \in \mathbb{C}^*} \\ \downarrow \text{res} & & \downarrow \beta \\ \text{Rep}(\mathcal{U}_q(\mathfrak{g})) & \xrightarrow{\chi} & \mathbb{Z}[y_i^\pm]_{i \in I} \end{array}$$

where β is the ring homomorphism such that $\beta(Y_{i,a}) = y_i$ ($i \in I, a \in \mathbb{C}^*$).

Example:

$$\begin{aligned} \chi_q(M_{aq^2} \otimes M_a) &= (Y_{aq^3} + Y_{aq}^{-1})(Y_{aq} + Y_{aq^{-1}}^{-1}) \\ &= Y_{aq^3}Y_{aq} + Y_{aq^3}Y_{aq^{-1}}^{-1} + Y_{aq}^{-1}Y_{aq^{-1}}^{-1} + 1 = \chi_q(N) + \chi_q(\text{trivial module}) \end{aligned}$$

where N is the simple module with Drinfeld polynomial $P(u) = (1 - aq^3u)(1 - aq^1u)$. In particular with $\dim(N) = 3$.

Note that $M_{aq^2} \otimes M_a \neq N \oplus \text{triv.}$ is not semi-simple. In fact we have an exact sequence:

$$0 \rightarrow N \rightarrow M_{aq^2} \otimes M_a \rightarrow \text{trivial module} \rightarrow 0$$

Example:

$$\chi_q(M_a \otimes M_a) = Y_a^2 + 2Y_aY_{aq^2}^{-1} + Y_{aq^2}^{-1}$$

In particular $M_a \otimes M_a$ is the simple module with Drinfeld polynomial $P(u) = (1 - au)^2$.

Definition 3. For $i \in I, a \in \mathbb{C}^*$ we note $V_{i,a}$ the simple module with Drinfeld polynomials $P_j(u) = \delta_{i,j}(1 - ua)$. Those simple modules are called *fundamental representations*.

We note $X_{i,a} = \chi_q(V_{i,a}) \in \mathcal{Y}$.

Corollary 1. (Frenkel, Reshetikhin 98) The ring $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ is commutative and isomorphic to $\mathbb{Z}[X_{i,a}]_{i \in I, a \in \mathbb{C}^*}$.

We say that $m \in \mathcal{Y}$ is a dominant monomial if it is of the form $m = \prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{u_{i,a}(m)}$ with $u_{i,a}(m) \geq 0$.

For m a dominant monomial we note $M_m \in \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$ the module $\bigotimes_{i \in I, a \in \mathbb{C}^*} V_{i,a}^{\otimes u_{i,a}(m)}$. It is called a standard module and his q -character is $\chi_q(M_m) = \prod_{i \in I, a \in \mathbb{C}^*} X_{i,a}^{u_{i,a}(m)}$.

6. NAKAJIMA'S q, t -CHARACTERS

In the ADE -case Nakajima defined a $\mathbb{Z}[t^\pm]$ -linear map :

$$\chi_{q,t} : \text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) \otimes_{\mathbb{Z}} \mathbb{Z}[t^\pm] \rightarrow \mathcal{Y}_t = \mathbb{Z}[Y_{i,a}^\pm, t^\pm]_{i \in I, a \in \mathbb{C}^*}$$

such that $(\chi_{q,t})_{t=1} = \chi_q$ but $\chi_{q,t} \neq \chi_q$. In particular it leads to the construction of :

- a quantization of $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}}))$
- an involution of $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) \otimes \mathbb{Z}[t^\pm]$
- canonical invariant basis of $\text{Rep}(\mathcal{U}_q(\hat{\mathfrak{g}})) \otimes \mathbb{Z}[t^\pm]$
- analogs of Kazhdan-Lusztig polynomials

Nakajima gave a combinatorial axiomatic definition of q, t -characters, but the existence is non-trivial and is proved with the geometric theory of quiver varieties which holds only in the ADE -case.

In the preprint math.QA/0212257 we propose a construction of $\chi_{q,t}$ without quiver varieties; in particular we extend the applications to the non-simply-laced case. In the following we give a sketch of the construction.

7. ALGEBRAIC CONSTRUCTION OF q, t -CHARACTERS IN THE GENERAL CASE

We set $\text{Rep} = \mathbb{Z}[X_{i,l}]_{i \in I, l \in \mathbb{Z}}$ and $\text{Rep}_t = \text{Rep} \otimes \mathbb{Z}[t^\pm]$

7.1. **Quantization of \mathcal{Y} .** Let Z_q be the \mathbb{C} -algebra defined by generators $a_i[m]$ ($i \in I, m \in \mathbb{Z} - \{0\}$), central elements c_r ($r > 0$) and relations ($i, j \in I, m, r \in \mathbb{Z} - \{0\}$):

$$[a_i[m], a_j[r]] = \delta_{m,-r}(q^m - q^{-m})B_{i,j}(q^m)c_{|m|}$$

For $j \in I, m \in \mathbb{Z}$ we set:

$$y_j[m] = \sum_{i \in I} \tilde{C}_{i,j}(q^m)a_i[m] \in Z_q$$

Consider the \mathbb{C} -algebra $Z_{q,h} = Z_q[[h]]$ and:

$$\tilde{Y}_{i,l} = \exp\left(\sum_{m>0} h^m y_i[m]q^{lm}\right)\exp\left(\sum_{m>0} h^m y_i[-m]q^{-lm}\right) \in Z_{q,h}$$

For $R \in \mathbb{Z}((q^{-1}))$, introduce:

$$t_R = \exp\left(\sum_{m>0} h^{2m} R(q^m)c_m\right) \in Z_{q,h}$$

We note $\mathcal{Y}_u \subset Z_{q,h}$ the subalgebra generated by the $t_R, \tilde{Y}_{i,l}^\pm$ ($i \in I, l \in \mathbb{Z}, R \in \mathbb{Z}((q^{-1}))$).

Definition 4. $\tilde{\mathcal{Y}}_t$ (resp. $Z_{q,t}$) is the quotient of \mathcal{Y}_u (resp. $Z_{q,h}$) by the relations $t_R = t_{R_0}$.

We note $t = t_0$. In particular in $\tilde{\mathcal{Y}}_t$ we have $t_R = t^{R_0}$ and $\tilde{\mathcal{Y}}_t$ is a $\mathbb{Z}[t^\pm]$ -algebra. We have defined a quantization of \mathcal{Y} :

Proposition 1. We have an isomorphism of $\mathbb{Z}[t^\pm]$ -vector space $\tilde{\mathcal{Y}}_t \simeq \mathcal{Y} \otimes \mathbb{Z}[t^\pm]$ and $\tilde{\mathcal{Y}}_t/(t-1) \simeq \mathcal{Y}$.

Example: In the sl_2 -case, $\tilde{\mathcal{Y}}_t$ is defined by generators t^\pm, \tilde{Y}_l^\pm ($l \in \mathbb{Z}$) and relations:

$$\tilde{Y}_l \tilde{Y}_k = t^s \tilde{Y}_k \tilde{Y}_l$$

where:

$$\begin{aligned} s &= 0 \text{ if } l - k = 1 + 2r, r \in \mathbb{Z} \\ s &= 2(-1)^r \text{ if } l - k = 2r, r > 0 \\ s &= 2(-1)^{r+1} \text{ if } l - k = 2r, r < 0 \\ s &= 0 \text{ if } l = k \end{aligned}$$

7.2. **Deformed screening operators.** Frenkel-Reshetikhin-Mukhin have shown $\text{Im}(\chi_q) = \bigcap_{i \in I} \text{Ker}(S_i)$

where the S_i are screening operators. We will use a t -version of this property. Introduce the screening currents:

$$\tilde{S}_{i,t} = \exp\left(\sum_{m>0} h^m \frac{a_i[m]}{q_i^m - q_i^{-m}} q^{lm}\right)\exp\left(\sum_{m>0} h^m \frac{a_i[-m]}{q_i^{-m} - q_i^m} q^{-lm}\right) \in Z_{q,t}$$

Definition 5. The i^{th} t -screening operator is the map $\tilde{S}_{i,t} : \tilde{\mathcal{Y}}_t \rightarrow Z_{q,t}$ defined by:

$$\tilde{S}_{i,t}(\lambda) = \frac{1}{t^2 - 1} \sum_{l \in \mathbb{Z}} [\tilde{S}_{i,t}, \lambda]$$

We note $\tilde{\mathfrak{K}}_t = \bigcap_{i \in I} \text{Ker}(\tilde{S}_{i,t})$. It is a subalgebra of $\tilde{\mathcal{Y}}_t$.

Theorem 4. For m a dominant monomial there a unique $\tilde{F}_t(m)$ in a completion of $\tilde{\mathfrak{K}}_t$ such that m is the unique dominant monomial of $\tilde{F}_t(m)$. Moreover it is given by a t -analog of Frenkel-Mukhin algorithm.

The proof of the existence is proved by showing that a t -analog of the algorithm is well-defined. See examples in the annexe.

7.3. Definition of q, t -characters.

Definition 6. *The morphism of q, t -characters is the map $\chi_{q,t} : \text{Rep}_t \rightarrow \tilde{\mathcal{Y}}_t^\infty$ which is $\mathbb{Z}[t^\pm]$ -linear and ($u_{i,l} \geq 0$):*

$$\chi_{q,t} \left(\prod_{i \in I, l \in \mathbb{Z}} X_{i,l}^{u_{i,l}} \right) = \prod_{l \in \mathbb{Z}} \tilde{F}_t(Y_{i,l})^{u_{i,l}}$$

Example:

Theorem 5. *We have $(\chi_{q,t})_{t=1} = \chi_q$. In particular the map $\chi_{q,t}$ is injective. In the ADE-case it is the morphism of Nakajima.*

7.4. Consequences. As $\chi_{q,t}$ is injective, the quantization of \mathcal{Y} leads to a quantization of Rep and the involution $c_r \mapsto c_r^{-1}$ of $Z_{q,h}$ leads to an involution of Rep_t .

Theorem 6. *For m a dominant monomial there is a unique $\tilde{L}_t(m) \in \text{Im}(\chi_{q,t})$ such that:*

$$\begin{aligned} \overline{\tilde{L}_t(m)} &= \tilde{L}_t(m) \\ \chi_{q,t}(M_m) &= \tilde{L}_t(m) + \sum_{m' < m, m' \text{ dominant}} P_{m',m}(t) \tilde{L}_t(m') \end{aligned}$$


where $P_{m',m}(t) \in t^{-1}\mathbb{Z}[t^{-1}]$.

Conjecture 1. *For m a dominant monomial, the image $(\tilde{L}_t(m))_{t=1}$ of $\tilde{L}_t(m)$ in \mathcal{Y} is $\chi_q(V_m)$ where V_m is the simple module of Drinfeld-polynomials associated to m .*


In the ADE-case the conjecture 1 is a consequence of Nakajima's geometric theory.

8. ANNEXE

8.1. We suppose $\mathfrak{g} = G_2$. The Cartan matrix is $C = \begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$ and $r_1 = 1$, $r_2 = 3$. The tree of the second fundamental representation is:



8.2. We suppose $\mathfrak{g} = A_2$. The Cartan matrix is $C = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ and $r_1 = 1$, $r_2 = 1$. The tree of $\tilde{F}_t(Y_{1,0}Y_{1,2}^2)$ is:



DAVID HERNANDEZ: ÉCOLE NORMALE SUPÉRIEURE - DMA, 45, RUE D'ULM F-75230 PARIS, CEDEX 05 FRANCE

E-mail address: David.Hernandez@ens.fr, URL: <http://www.dma.ens.fr/~dhernand>